Index to Volume 122

The Auk 122(4):1314–1341, 2005
© The American Ornithologists' Union, 2005.
Printed in USA.

Compiled by Todd J. Underwood and Robyn M. Underwood

The index is confined primarily to new information. New specific names are printed in **boldface** type.

100 Years Ago in the American Ornithologists' Union, 379–380, 722–723, 1024–1025, 1310–1311
Abel, Becky, see Mehlman, David W., et al.
Able, Kenneth P., see Holberton, Rebecca L., and –
abundance, 15–25, 338–346, 1249–1260; food –, 783–792, 1225–1238; fruit –, 783–792; prey –, 887–901
Acanthiza reguloides, 1097–1111
Acanthorhynchus tenuirostris, 1097–1111
Accipiter, 108–120
A. badius, 108–120
A. cooperii, 1037–1048, 1130–1148
A. striatus, 1130–1148
A. trivirgatus, 108–120
A. virgatus, 108–120
acoustic, – adaptation hypothesis, 231–245; – signals, 231–245
Acridotheres cristatellus, 1026–1031
A. javanicus, 1026–1031
Acrocephalus arundinaceus, 1–14, 530–543, 558–565, 1086–1096
A. bistrigiceps, 1086–1096
A. melanopogon, 835–842
A. newtoni, 1271–1280
A. paludicola, 558–565
A. schoenobaenus, 558–565
A. scirpaceus, 530–543, 558–565, 835–842, 1291–1292
A. stentoreus, 558–565
Actitis macularius, 1130–1148
Adam, Aileen, see Gorman, Helen E., et al.
adaptation, local –, 803–818; social –, 803–818
adaptive radiation, 121–143
Aechmophorus occidentalis, 1130–1148
aerodynamics, 872–886
Africa, 1271–1280
age differences, 1070–1085; – ratios, 1070–1085
age-class, 915–921
Agelaius phoeniceus, 57–70, 82–93, 1037–1048, 1130–1148
aggression, 530–543; interspecific –, 1097–1111
Aimophila ruficeps, 571–592
alarm calls, 108–120
Alaudidae, 733–746
Albatross, 1182–1190; Black-browed, 1182–1190; Gray-headed, 1182–1190; Light-Mantled Sooty, 773–782; Sooty, 773–782; Wandering, 1182–1190
Alca impennis, 971–979
A. torda, 57–70, 773–782, 887–901
Alcedinidae, 1271–1280
Alcedo, 1271–1280
A. atthis, 1271–1280
A. cristata, 1271–1280
A. cyanopecta, 1271–1280
A. c. cyanopecta, 1271–1280
A. c. nigrirosta, 1271–1280
A. leucogaster, 1271–1280
A. menninting, 1271–1280
A. quadribrachys, 1271–1280
A. semitorquata, 1271–1280
A. vintsioides, 1271–1280
alcid, 887–901
Alectroenas, 1271–1280
Alectura lathami, 1239–1248
Alisauskas, Ray T., see Kellett, Dana K., et al.
Altshuler, Douglas L., see Stiles, F. Gary, et al.
Amadon, Dean, see Delacour, Jean, et al.
Amazilia, 593–607
A. saucerottei, 187–204
Amazon, Orange-winged, 292–305
Amazona amazonica, 292–305
American Museum of Natural History, 389–402
Ammodramus, 94–107
A. bairdii, 349–350
A. caudacutus, 94–107
A. c. caudacutus, 94–107
A. c. diverus, 94–107
A. henslowii, 618–636, 661–672
A. nelsoni alterus, 94–107
A. n. nelsoni, 94–107
A. n. subvirgatus, 94–107
A. savannarum, 618–636
Amphispiza belli, 571–592
Anas acuta, 1130–1148
A. americana, 1130–1148
A. clypeata, 1130–1148
A. crecca, 1130–1148
A. discors, 1130–1148
A. fulvigula, 523–529, 949–965
A. platyrhynchos, 255–267, 509–522, 523–529, 558–565, 949–965, 1130–1148
A. p. domesticus, 523–529
A. rubripes, 523–529, 949–965

Index to Volume 122

The Auk 122(4):1314–1341, 2005
© The American Ornithologists' Union, 2005.
Printed in USA.

Compiled by Todd J. Underwood and Robyn M. Underwood

The index is confined primarily to new information. New specific names are printed in **boldface** type.

100 Years Ago in the American Ornithologists' Union, 379–380, 722–723, 1024–1025, 1310–1311
Abel, Becky, see Mehlman, David W., et al.
Able, Kenneth P., see Holberton, Rebecca L., and –
abundance, 15–25, 338–346, 1249–1260; food –, 783–792, 1225–1238; fruit –, 783–792; prey –, 887–901
Acanthiza reguloides, 1097–1111
Acanthorhynchus tenuirostris, 1097–1111
Accipiter, 108–120
A. badius, 108–120
A. cooperii, 1037–1048, 1130–1148
A. striatus, 1130–1148
A. trivirgatus, 108–120
A. virgatus, 108–120
acoustic, – adaptation hypothesis, 231–245; – signals, 231–245
Acridotheres cristatellus, 1026–1031
A. javanicus, 1026–1031
Acrocephalus arundinaceus, 1–14, 530–543, 558–565, 1086–1096
A. bistrigiceps, 1086–1096
A. melanopogon, 835–842
A. newtoni, 1271–1280
A. paludicola, 558–565
A. schoenobaenus, 558–565
A. scirpaceus, 530–543, 558–565, 835–842, 1291–1292
A. stentoreus, 558–565
Actitis macularius, 1130–1148
Adam, Aileen, see Gorman, Helen E., et al.
adaptation, local –, 803–818; social –, 803–818
adaptive radiation, 121–143
Aechmophorus occidentalis, 1130–1148
aerodynamics, 872–886
Africa, 1271–1280
age differences, 1070–1085; – ratios, 1070–1085
age-class, 915–921
Agelaius phoeniceus, 57–70, 82–93, 1037–1048, 1130–1148
aggression, 530–543; interspecific –, 1097–1111
Aimophila ruficeps, 571–592
alarm calls, 108–120
Alaudidae, 733–746
Albatross, 1182–1190; Black-browed, 1182–1190; Gray-headed, 1182–1190; Light-Mantled Sooty, 773–782; Sooty, 773–782; Wandering, 1182–1190
Alca impennis, 971–979
A. torda, 57–70, 773–782, 887–901
Alcedinidae, 1271–1280
Alcedo, 1271–1280
A. atthis, 1271–1280
A. cristata, 1271–1280
A. cyanopecta, 1271–1280
A. c. cyanopecta, 1271–1280
A. c. nigrirosta, 1271–1280
A. leucogaster, 1271–1280
A. menninting, 1271–1280
A. quadribrachys, 1271–1280
A. semitorquata, 1271–1280
A. vintsioides, 1271–1280
alcid, 887–901
Alectroenas, 1271–1280
Alectura lathami, 1239–1248
Alisauskas, Ray T., see Kellett, Dana K., et al.
Altshuler, Douglas L., see Stiles, F. Gary, et al.
Amadon, Dean, see Delacour, Jean, et al.
Amazilia, 593–607
A. saucerottei, 187–204
Amazon, Orange-winged, 292–305
Amazona amazonica, 292–305
American Museum of Natural History, 389–402
Ammodramus, 94–107
A. bairdii, 349–350
A. caudacutus, 94–107
A. c. caudacutus, 94–107
A. c. diverus, 94–107
A. henslowii, 618–636, 661–672
A. nelsoni alterus, 94–107
A. n. nelsoni, 94–107
A. n. subvirgatus, 94–107
A. savannarum, 618–636
Amphispiza belli, 571–592
Anas acuta, 1130–1148
A. americana, 1130–1148
A. clypeata, 1130–1148
A. crecca, 1130–1148
A. discors, 1130–1148
A. fulvigula, 523–529, 949–965
A. platyrhynchos, 255–267, 509–522, 523–529, 558–565, 949–965, 1130–1148
A. p. domesticus, 523–529
A. rubripes, 523–529, 949–965

A. strepera, 1130–1148
A. superciliosa, 949–965
A. wyvilliana, 949–965
A. zonorhyncha, 949–965
Anatidae, 949–965
Anatinae, 313–318
Anderson, Kurt E., et al., Large-scale movement patterns between song dialects in Brown-headed Cowbirds (*Molothrus ater*), 803–818
Andes, 445–463
Andigena cucullata, 1191–1209
animal model, 1123–1129
annual cycle energetics, 922–937
Anser albifrons, 1130–1148
Anseriformes, 566–570, 971–979, 1130–1148
Anserinae, 313–318
Antbird, 403–413, 433–444, 673–683; Bicolored, 843–852; Spotted, 843–852
Anthochaera curunculata, 1097–1111
Anthoscopus, 121–143
A. minutus, 121–143
anthropogenic noise, 231–245
Anthus pratensis, 530–543
A. spragueii, 349–350
anticipation hypothesis, 509–522
Antpitta, 673–683; Moustached, 389–402; Peruvian, 389–402; Streak-chested, 843–852; Watkin's, 389–402
Antshrike, Variable, 403–413, 414–432, 433–444
Ant-Tanager, Red-throated, 783–792
antthrushes, 673–683
Antwren, Checker-throated, 843–852
Aphelocoma californica, 464–478, 571–592
A. coerulescens, 464–478, 571–592
A. ultramarina, 571–592, 1130–1148
Aphrastura, 158–174
A. spinicauda, 158–174
Apodiformes, 971–979, 1049–1054
Araçari, Lettered, 1191–1209
Aramus guarauna, 26–35
Aratinga, 292–305
A. acuticaudata, 292–305
A. aurea, 292–305
A. auricapillus, 292–305
A. a. auricapillus, 292–305
A. a. aurifrons, 292–305
A. brevipes, 292–305
A. cactorum, 292–305
A. canicularis, 292–305
A. chloroptera, 292–305
A. erythrogenys, 292–305
A. euops, 292–305
A. finschi, 292–305
A. holochlora, 292–305
A. jandaya, 292–305
A. leucophthalma, 292–305
A. mitrata, 292–305
A. nana, 292–305
A. pertinax, 292–305
A. pintoi, 292–305
A. rubritorquis, 292–305
A. solstitialis, 292–305
A. wagleri, 292–305
A. weddellii, 292–305
Archilochus alexandri, 187–204, 872–886, 938–948, 1097–1111
A. colubris, 872–886, 938–948
archive, digital –, 984–987
Arctic species clade, 684–688
Arctic-nesting, – birds, 479–496; – species, 313–318
Ardea alba, 255–267
A. herodias, 1130–1148
area-restricted search, 36–49
area-sensitive, 618–636
Arenaria interpres, 1149–1160
Arendt, Wayne J., review by, 1016–1018
Armstrong, Jessica L., see Cristol, Daniel A., et al.
Arsenault, David P., et al., Mark–recapture and DNA fingerprinting data reveal high breeding-site fidelity, low natal philopatry, and low levels of genetic population differentiation in Flammulated Owls (*Otus flammeolus*), 329–337
artificial, – insemination, 1161–1168; – nests, 36–49, 843–852
Asio capensis, 1271–1280
A. flammeus, 1130–1148
asities, 1271–1280
aspect ratio, 872–886
asymmetry, 1123–1129
Athene cunicularia, 464–478
A. c. floridana, 464–478
A. c. hypugaea, 464–478
attenuate, 231–245
Audubon, 347–348
Auk, Great, 971–979
Auklet, Rhinoceros, 887–901
Aulacorhynchus derbianus, 1055–1063
A. prasinus, 1191–1209
Auriparus, 121–143
A. flaviceps, 121–143
Australia, 1097–1111
autumn migration timing, 1070–1085
availability, food –, 144–157, 902–914; resource –, 544–557, 1281–1290
avian knowledge network, 15–25
avifauna, 1271–1280
Aythya affinis, 1130–1148
A. americana, 1130–1148
A. collaris, 1130–1148
A. valisineria, 1037–1048, 1130–1148

Babbler, Orange-billed, 108–120
Bacillus licheniformis, 222–230
badge size, 835–842
Baeolophus, 121–143, 747–758
B. atricristatus, 747–758

B. bicolor, 108–120, 121–143
B. inornatus, 121–143, 747–758
B. ridgwayi, 747–758
B. wollweberi, 121–143
Baillonius bailloni, 1055–1063, 1191–1209
Baker, Allan J., see Given, Andrew D., et al.
Baker, Myron C., see Gammon, David E., et al.
Balda, Russell P., 359–361
Bananaquit, 187–204
band, – recovery, 479–496; – wear, 773–782
banding, 71–81
Bangsia melanochlamys, 445–463
Barber, Brian R., see Martínez Gómez, Juan E., et al.
barbets, 1049–1054, 1055–1063
Barro Colorado Island, 389–402
Bart, Jonathan, Monitoring the abundance of bird populations, 15–25
Bartholomew, George A., see Cade, Tom J., et al.
Bartlett, Terri L., et al., Division of labor: Incubation and biparental care in House Sparrows (*Passer domesticus*), 835–842
Battley, Phil F., and Piersma, Theunis, Body composition and flight ranges of Bar-tailed Godwits (*Limosa lapponica baueri*) from New Zealand, 922–937
Beardmore, C. J., see Rich, T. D., et al.
Bednarz, James C., see Bollmer, Jennifer L., et al.
Bee-eater, 733–746; Little, 1271–1280; Red-throated, 1271–1280
behavior, 1–14, 158–174, 347–348; begging –, 1112–1122; female –, 57–70; flight –, 872–886; flight-calling –, 733–746; flocking –, 1249–1260; foraging –, 36–49, 187–204; parental –, 1239–1248; post-fledging –, 205–221; ranging –, 1249–1260; renesting –, 1000–1001; reproductive –, 347–348; sex-specific –, 347–348; vocal –, 175–186
behavioral plasticity, 566–570
Bell, Douglas A., see Gay, Laurène, et al.
Bellrose, Frank C., 1916–2005, In Memoriam, 1298–1300
Benson, Anna-Marie, and Winker, Kevin, Fat-deposition strategies among high-latitude passerine migrants, 544–557
Bergier, Patrick, see Thévenot, Michel, et al.
Berlanga, H., see Rich, T. D., et al.
berries, 144–157
Bidwell, Mark T., and Dawson, Russell D., Calcium availability limits reproductive output of Tree Swallows (*Tachycineta bicolor*) in a nonacidified landscape, 246–254
bill, – morphology, 187–204; – shape, 1191–1209,
binomial distribution, 1161–1168
bioaccumulation, 255–267
bioacoustics, 984–987
biodiversity, 966–971; – information system, 987–990
biogeography, 50–56, 121–143, 158–174, 268–269, 389–402, 414–432, 445–463, 1037–1048
bioindicators, 966–971
bioinformatics, 979–984
biological species concept, 50–56, 403–413, 433–444
biomonitors, 990–994
biparental care, 835–842
Birdlife International, Tracking Ocean Wanderers: The Global Distribution of Albatrosses and Petrels (rev.), 1307–1308
birdsong, 853–871
Birks, Sharon, see Edwards, Scott V., et al.
Bishop, 57–70; Red, 57–70
Bittern, 26–35; American, 26–35, 1130–1148; Least, 26–35
Blackbird, 733–746; Brewer's, 82–93; Eurasian, 530–543, 558–565; European, 733–746; Red-winged, 57–70, 82–93, 1037–1048, 1130–1148; Yellow-headed, 1130–1148
Blackcap, 530–543, 733–746
Blancher, P. J., see Rich, T. D., et al.
Blancher, Peter J., see Dunn, Erica H., et al.
blood, – chemistry, 1261–1270; – parasites, 558–565
Bluebird, Eastern, 1086–1096; Mountain, 1130–1148
Bluethroat, 71–81
Bobolink, 618–636, 701–705, 733–746, 1130–1148
Bock, Walter J., In Memoriam: Ernst Mayr, 1904–2005, 1005–1007
body composition, 922–937; – mass, 313–318; – size, 922–937
Bollmer, Jennifer L., et al., Population genetics of the Galápagos Hawk (*Buteo galapagoensis*): Genetic monomorphism within isolated populations, 1210–1224
boluses, 1182–1190
Bombycilla cedrorum, 1070–1085
bones, 971–979
Borgia, Gerald, review by, 718–721
Botaurus lentiginosus, 26–35, 1130–1148
Brachypteraciidae, 1049–1054
Brachyramphus marmoratus, 36–49, 990–994
Bradbury, Jack W., see Gaunt, Sandra L. L., et al.
Bradstreet, M. S. W., see Rich, T. D., et al.
Branta canadensis, 144–157, 523–529, 1037–1048, 1130–1148
Branta c. interior, 144–157
B. c. maxima, 144–157
B. hutchinsii minima, 144–157
B. leucopsis, 479–496
B. sandvicensis, 329–337, 1026–1031
Braun, Michael J., see Bronson, C. L., et al.
Brawn, Jeffrey D., see Robinson, W. Douglas, et al.
Brazil, 292–305
Breeding Bird Survey, 338–346
breeding, 637–647; – colonies, 773–782; – phenology, 571–592; – season, 887–901
Breton, André R., et al., Adult survival estimates from two Atlantic Puffin (*Fratercula arctica*) colonies in the Gulf of Maine, 773–782
Brewster Memorial Award, 2004: Russell P. Balda and Alan C. Kamil, 359–361

brightness, 793–802
Bristle-Tyrant, Chapman's, 389–402
broadbill, 1271–1280
Brodin, Anders, Hippocampal volume does not correlate with food-hoarding rates in the Black-capped Chickadee (*Poecile atricapillus*) and Willow Tit (*Parus montanus*), 819–828
Bronson, C. L., et al., Reproductive success across the Black-capped Chickadee (*Poecile atricapillus*) and Carolina Chickadee (*P. carolinensis*) hybrid zone in Ohio, 759–772
brood, – loss, 479–496; – parasite density, 648–660; – parasitism, 280–291, 530–543, 648–660; – patch, 835–842; – reduction, 509–522
brood-rearing, 144–157
broods, successful –, 1001–1003
Brooks, Daniel M., review by, 1018–1019
Brotogeris versicolurus, 292–305
Brumfield, Robb T., Mitochondrial variation in Bolivian populations of the Variable Antshrike (*Thamnophilus caerulescens*), 414–432
Brumfield, Robb T., see Edwards, Scott V., et al.
Brumfield, Robb T., see Isler, Morton L., et al.
Brush-turkeys, Australian, 1239–1248
Bucephala albeola, 1130–1148
B. clangula, 313–318, 1130–1148
Budgerigar, 733–746
Budney, Gregory F., see Gaunt, Sandra L. L., et al.
Bufflehead, 1130–1148
Bulbul, 1271–1280; Madagascar, 1271–1280; Yellow-browed, 108–120
Bullfinch, Lesser Antillean, 187–204
Bunting, 747–758, 938–948; Corn, 853–871; Indigo, 280–291, 733–746, 938–948, 1086–1096; Lazuli, 938–948, 1070–1085; Ortolan, 733–746; Painted, 938–948, 1070–1085; Rustic, 733–746
Burhinus bistriatus, 1026–1031
Butcher, G. S., see Rich, T. D., et al.
Buteo, 1210–1224
B. buteo, 1026–1031
B. galapagoensis, 1210–1224
B. jamaicensis, 1130–1148
B. lagopus, 1130–1148
B. swainsonii, 1210–1224
Butorides virescens, 26–35

cache, 819–828
Cacicus cela, 593–607
Caciques, Yellow-rumped, 593–607
Cactus-Finch, Large, 571–592
Cade, Tom J., et al., In Memoriam: Thomas Raymond Howell, 1924–2004, 1008–1010
Cadena, Carlos Daniel, see Cuervo, Andrés M., et al.
Cadieux, Marie-Christine, et al., Feeding ecology of Canada Geese (*Branta canadensis interior*) in sub-Arctic inland tundra during brood-rearing, 144–157
Calcarius ornatus, 701–705
calcium limitation, 246–254
Calidris alpina, 1037–1048, 1249–1260
C. canutus, 922–937, 1249–1260
C. tenuirostris, 922–937
California coastal scrub, 571–592
call-broadcast surveys, 26–35
calling, reciprocal –, 733–746
calls, alarm –, 108–120; flight –, 733–746
Caloenas nicobarica, 1003–1004
Calonectris diomedea, 255–267
Calypte anna, 187–204, 872–886
Campephilus principalis, 389–402, 984–987
Campethera abingoni, 1055–1063
C. cailliautii, 1055–1063
C. nivosa, 1055–1063
Campylopterus curvipennis, 593–607
C. excellens, 593–607
Canary, Common, 1–14
candling, 661–672
Cannon, Michelle D., see Bollmer, Jennifer L., et al.
Canvasback, 1037–1048, 1130–1148
Capido squamatus, 1055–1063
capital breeders, 1225–1238
Capitonidae, 1055–1063
Capitonides, 1055–1063
Capparella, Angelo P., see Cheviron, Z. A., et al.
Caprimulgiformes, 971–979, 1130–1148
Caprimulgus, 1271–1280
capture–mark–recapture, 319–328
capture–mark–resight/recapture, 773–782
capture–recapture, 637–647
Cardinal, Northern, 222–230
cardinalid, 938–948
Cardinalis cardinalis, 222–230
cardueline, 733–746
Carduelis pinus, 1070–1085
Carib, Purple-throated, 187–204
Carlisle, Jay D., et al., Molt strategies and age differences in migration timing among autumn landbird migrants in southwestern Idaho, 1070–1085
carotenoid, 793–802
Carpodacus cassinii, 1070–1085
C. mexicanus, 403–413
C. purpureus, 1130–1148
Catbird, Gray, 938–948, 1130–1148
Cathartes aura, 389–402
Catharus, 733–746
C. fuscescens, 733–746
C. guttatus, 689–700, 938–948, 1070–1085, 1130–1148
C. minimus, 544–557, 733–746
C. ustulatus, 544–557, 733–746, 938–948, 1037–1048, 1070–1085
Catoptrophorus semipalmatus, 1130–1148, 1249–1260
Causey, Douglas, and Trimble, Jeremiah, Old bones in new boxes: Osteology collections in the new millennium, 971–979
Centropus, 1271–1280

Cephalopoda, 1182–1190
Cercomacra parkeri, 445–463
Cercotrichas galactotes, 1169–1181
Cerorhinca monocerata, 887–901
Certhia americana, 50–56, 1070–1085, 1130–1148
Ceryle alcyon, 1130–1148
C. rudis, 1271–1280
Cerylidae, 1271–1280
Ceycoides, 1271–1280
C. madagascariensis, 1271–1280
Ceyx, 1271–1280
C. erithacus, 1271–1280
C. lecontei, 1271–1280
C. lepidus, 1271–1280
C. madagascariensis, 1271–1280
C. melanurus melanurus, 1271–1280
C. m. mindanensis, 1271–1280
C. m. samarensis, 1271–1280
C. picta, 1271–1280
C. rufidorsum, 1271–1280
Chaetura champani, 389–402
Chamaeza, 673–683
C. campanisona, 673–683
Chapman Memorial Fund, 389–402
Chapman, Frank M., 389–402
character, – evolution, 1191–1209; – reconstruction, 1191–1209
Charadriiformes, 971–979, 1130–1148
Charadrius melodus, 15–25
C. montanus, 1037–1048
C. vociferus, 1130–1148
C. wilsonia, 1249–1260
Chase, Mary K., et al., Effects of weather and population density on reproductive success and population dynamics in a Song Sparrow (*Melospiza melodia*) population: A long-term study, 571–592
Chat, White-fronted, 280–291; Yellow-breasted, 938–948
Chen caerulescens, 313–318, 1130–1148
C. c. atlantica, 144–157, 479–496
C. c. caerulescens, 144–157, 479–496
C. canagica, 479–496
C. rossii, 313–318
Cheviron, Z. A., et al., Molecular phylogenetic relationships among the *Geositta* miners (Furnariidae) and biogeographic implications for avian speciation in Fuego-Patagonia, 158–174
Chickadee, 121–143, 747–758; Black-capped, 108–120, 689–700, 747–758, 759–772, 819–828, 853–871, 1070–1085; Boreal, 747–758; Carolina, 747–758, 759–772; Chestnut-backed, 82–93; Gray-headed, 747–758; Mexican, 747–758; Mountain, 747–758, 853–871, 1070–1085
chicken, 158–174, 414–432, 835–842, 843–852, 949–965
chick-rearing period, 1182–1190
Chiffchaff, 733–746, 1291–1292
Chlidonias niger, 1130–1148
Chloroceryle americana, 1271–1280
choice, female –, 82–93; mate –, 1–14, 71–81, 82–93, 1169–1181
Chondestes grammacus, 1070–1085
Chordeiles minor, 1130–1148
Christmas Bird Count, 205–221, 338–346, 389–402
chroma, 793–802
Chrysocolaptes lucidus, 1055–1063
Cicero, Carla, see Peterson, A. Townsend, et al.
Cichoń, Mariusz, see Dubiec, Anna, et al.
Ciconiiformes, 255–267, 971–979, 1130–1148
Cimprich, David, see Mehlman, David W., et al.
Cinclodes, 158–174
C. excelsior, 158–174
Cinclus cinclus, 637–647
Circus aeruginosus, 1026–1031
C. cyaneus, 1026–1031, 1130–1148, 1291–1292
C. pygargus, 1291–1292
C. spilonotus, 1026–1031
Cisticola, 1271–1280
Cistothorus palustris, 26–35, 1130–1148
C. platensis, 50–56
citizen science, 15–25, 338–346
Clamator jacobinus, 733–746
Clangula hyemalis, 313–318
classification, 403–413
climate, 637–647; – change, 1037–1048, 1130–1148
clinal variation, 414–432, 433–444
Clotfelter, Ethan D., see Sampieri, Sara J., and –
cloud forest, 1064–1069
clutch, – initiation date, 246–254; – size, 246–254, 648–660, 1003–1004, 1225–1238
CMR, 773–782
Coccyzus erythropthalmus, 733–746
Cock-of-the-Rock, 793–802
Coereba flaveola, 187–204, 1026–1031
Coerebidae, 1026–1031
coevolution, 530–543
Colaptes auratus, 1055–1063, 1070–1085, 1130–1148
Coliidae, 1055–1063
Collared-Dove, Eurasian, 530–543
collections, bird –, 966–971, 990–994; eggshell –, 994–999; frozen –, 979–984; genetic resources –, 979–984; museum –, 938–948, 971–979, 987–990; osteology –, 971–979
Colombia, 445–463
colonial waterbirds, 464–478
colonies, 497–508; breeding –, 773–782
color, 793–802
coloration, 1191–1209
Columba livia, 530–543, 608–617, 733–746
C. l. domestica, 530–543
columbids, 1003–1004
Columbiformes, 971–979, 1130–1148
community ecology, 108–120
comparative function, 971–979
competition, 1112–1122; food –, 1055–1063

condition, 829–834; body –, 544–557, 558–565; female –, 509–522, 829–834, 1225–1238; maternal –, 1161–1168; nestling –, 1225–1238; nutritional –, 1261–1270; offspring –, 1112–1122; physiological –, 1261–1270
conditional-extrapair-strategy hypothesis, 57–70
connectivity, 464–478, 1149–1160, 1281–1290; habitat –, 853–871; migratory –, 1037–1048
Conopophaga, 673–683
C. ardesciaca, 673–683
C. aurita, 673–683
C. castaneiceps, 673–683
C. lineata, 673–683
C. melanogaster, 673–683
C. melanops, 673–683
C. peruviana, 673–683
C. roberti, 673–683
Conopophagidae, 673–683
conservation, 338–346, 733–746, 1281–1290
conspecific brood parasitism, 566–570
constraint hypothesis, 509–522
contact zones, 747–758
contaminants, 990–994; industrial –, 915–921
Contopus sordidulus, 82–93, 1070–1085
C. virens, 1070–1085
control region, 684–688
Conuropsis carolinensis, 389–402
Conway, Courtney J., and Gibbs, James P., Effectiveness of call-broadcast surveys for monitoring marsh birds, 26–35
Cooper, Robert J., and Warnell, Daniel B., review by, 1019–1020
cooperative polyandry, 1210–1224
Coot, American, 26–35, 566–570, 1130–1148
Copsychus albospecularis, 1271–1280
copulation, 82–93
Coracias abyssinica, 1271–1280
Coraciidae, 1049–1054
Coraciiformes, 971–979, 1130–1148
Coracina, 1271–1280
Coracopsis, 1271–1280
Cormorant, 1182–1190; Double-crested, 1130–1148
corticosterone, 1–14
Corvidae, 36–49, 121–143, 733–746, 819–828
Corvus brachyrhynchos, 36–49, 205–221
C. corax, 36–49
C. corone, 36–49
Corythornis, 1271–1280
C. cristata, 1271–1280
C. leconteі, 1271–1280
C. leucogaster, 1271–1280
C. madagascariensis, 1271–1280
C. picta, 1271–1280
C. vintsioides, 1271–1280
Coturnicops noveboracensis, 26–35
Coturnix, 843–852
C. japonica, 36–49, 1037–1048
Coua, 1271–1280
Coues Award, 2004: Jared Verner, 361–362
courtship, 82–93
Cowbird, Brown-headed, 82–93, 94–107, 530–543, 571–592, 618–636, 648–660, 803–818, 1130–1148
Coyne, Jerry A., and Orr, H. Allen, Speciation (rev.), 371–373
Crane, Sandhill, 1130–1148
Crax nigra, 971–979
Creatophora cinerea, 1271–1280
Creeper, Brown, 50–56, 1070–1085, 1130–1148
Cresswell, Kate A., see Xavier, J. C., et al.
Cristol, Daniel A., et al., Feather-degrading bacteria do not affect feathers on captive birds, 222–230
Cristol, Daniel A., see Keiser, Jeffrey T., et al.
croakers, channel-keel-billed, 1191–1209
Crochet, Pierre-André, see Gay, Laurène, et al.
crop milk, 1003–1004
Crossbill, 1123–1129, 1225–1238; European Red, 1123–1129; Himalayan Red, 1123–1129; Parrot, 1123–1129; Red, 1123–1129; Scottish, 1123–1129; White-winged, 1123–1129
Crow, American, 36–49, 205–221; Carrion, 36–49
Croxall, John P., see Xavier, J. C., et al.
cryptic species, 1291–1292
Cuckoo, 733–746, 1271–1280; Black-billed, 733–746; Common, 530–543, 1130–1148; Pied, 733–746
Cuculiformes, 971–979
Cuculus canorus, 530–543, 1130–1148
Cuervo, Andrés M., et al., *Scytalopus stilesi*, a new species of tapaculo (Rhinocryptidae) from the Cordillera Central of Colombia, 445–463
Cully, Jack F., Jr., see Jensen, William E., and –
cultural evolution, 853–871
Curlew, Long-billed, 15–25
Currasow, Great, 971–979
Currie, Phillip J., et al., Feathered Dragons: Studies on the Transition from Dinosaurs to Birds (rev.), 714–716
Curry, Robert L., Hybridization in chickadees: Much to learn from familiar birds, 747–758
Cyanistes, 121–143
C. caeruleus, 121–143, 747–758
C. cyanus, 121–143, 747–758
C. flavipectus, 747–758
Cyanocitta stelleri, 36–49, 571–592, 1070–1085
Cygnus columbianus, 1130–1148
cytochrome *b*, 121–143, 445–463, 673–683, 684–688
Czyrnyj, Ashleigh A., see Murphy-Klassen, Heather M., et al.

Dacelonidae, 1271–1280
Dantzker, Marc S., see Gaunt, Sandra L. L., et al.
Dawson, Russell D, see Bidwell, Mark T., and –
de Lima, Flávio César Thadeo, see Silveira, Luís Fábio, et al.
de Vries, Tjitte, see Bollmer, Jennifer L., et al.
defense, economics of –, 187–204; parental nest –, 701–705; resource –, 1097–1111

del Hoyo, J., see Delacour, Jean, et al.
Delacour, Jean, et al., Currasows and Related Birds (rev.), 1018–1019
Delichon, 121–143
D. urbicum, 637–647, 1055–1063
Delta Marsh, 1130–1148
Delta Waterfowl and Wetlands Research Station, 1130–1148
Demarest, D. W., see Rich, T. D., et al.
demographics, 464–478
demography, 306–312, 306–312, 637–647
Dendrocolaptidae, 673–683
Dendrocopos major, 1055–1063
D. medius, 1055–1063
D. minor, 1055–1063
Dendroica, 793–802, 1037–1048
D. caerulescens, 71–81, 306–312, 571–592, 793–802, 902–914, 1000–1001, 1001–1003, 1037–1048, 1086–1096
D. cerulea, 497–508
D. coronata, 71–81, 544–557, 733–746, 1070–1085, 1130–1148
D. c. coronata, 793–802, 938–948, 1037–1048
D. discolor, 280–291, 689–700
D. kirtlandii, 689–700
D. magnolia, 71–81, 793–802, 938–948, 1070–1085
D. nigrescens, 938–948
D. occidentalis, 938–948, 1070–1085
D. palmarum, 1130–1148
D. pensylvanica, 793–802
D. petechia, 530–543, 544–557, 1070–1085, 1130–1148
D. p. aestiva, 938–948
D. striata, 544–557, 689–700, 733–746
D. townsendi, 938–948, 1070–1085
D. virens, 938–948
Dendropicos fuscescens, 1055–1063
density, – dependence, 571–592; – independence, 571–592
Derham, William, 706–707
dermestid beetles, 971–979
detectability, 15–25
detection, – efficiency, 608–617; – probability, 26–35; – ratios, 15–25
deuterium, 1037–1048
development, early –, 1239–1248
developmental stress hypothesis, 1–14
DeVink, Jean-Michel A., et al., Effects of water salinity on growth and survival of Common Eider (*Somateria mollissima*) ducklings, 523–529
dialects, 593–607
Diamond, Antony W., see DeVink, Jean-Michel A., et al.
Diamond, Antony W., see Breton, André R., et al.
Dickcissel, 82–93, 648–660
Dickinson, Edward C., The Howard and Moore Complete Checklist of the Birds of the World, 3rd ed. (rev.), 712–714
Dicrurus, 1271–1280
D. paradiseus, 108–120
diet, 36–49, 144–157, 783–792, 887–901, 1182–1190, 1239–1248, 1249–1260, 1261–1270
differential leukocyte counts, 829–834
digital, – archive, 984–987; – sound recording, 984–987; – vouchers, 979–984
Dinopium javense, 1055–1063
Diomedia, 1182–1190
D. exulans, 1182–1190
Dipper, White-throated, 637–647
discrimination, 530–543
dispersal, 205–221, 329–337, 759–772, 803–818, 949–965, 1210–1224; natal –, 853–871
distress, 108–120
distributed generic information retrieval, 987–990
distribution, 338–346
diurnal studies, 1249–1260
diversity, genetic –, 94–107, 464–478, 1210–1224
division of labor, 835–842
DNA, – barcoding, 979–984; – fingerprinting, 329–337; – sequences, 121–143, 673–683, 979–984; – sequencing, 949–965
Dodo, 1003–1004
Dolichonyx oryzivorus, 618–636, 701–705, 733–746, 1130–1148
Doliornis remseni, 445–463
domestic fowl, 255–267
dominance, 1097–1111; – status, 759–772
Donacobius atricapilla, 1026–1031
Doran, Patrick J., see Jones, Jason, et al.
dormitory nests, 1169–1181
Dove, Brown-backed, 1026–1031; Gray-fronted, 1026–1031; Gray-headed, 1026–1031; Mourning, 1130–1148
Drake, Kiel L., see Kellett, Dana K., et al.
Drennan, Susan Roney, see Dunn, Erica H., et al.
Drongo, Greater Racket-tailed, 108–120
Drovetski, Sergei V., see Kulikova, Irina V., et al.
Dryocopus martius, 1055–1063
Dubiec, Anna, et al., Seasonal decline in leukocyte concentrations and reproductive output in female Great Tits (*Parus major*), 829–834
Duck, 313–318; American Black, 523–529, 949–965; Eastern Spot-billed, 949–965; Grey, 949–965; Hawaiian, 949–965; Long-tailed, 313–318; Mallard, 949–965; Mottled, 523–529, 949–965; Mountain, 523–529; Pekin, 523–529; Ring-necked, 1130–1148; Ruddy, 1130–1148
duckling, 523–529
Dudley, Robert, see Stiles, F. Gary, et al.
duet, 175–186
Dumetella carolinensis, 938–948, 1130–1148
Duncan, Charles, see Mehlman, David W., et al.
Dunlin, 1037–1048, 1249–1260
Dunn, E. H., see Rich, T. D., et al.
Dunn, Erica H., et al., Enhancing the scientific value of the Christmas Bird Count, 338–346
Dunn, Peter O., see Nooker, Jacqueline K., et al.

Dunnock, 57–70
Dyrcz, Andrzej, et al., Male reproductive success is correlated with blood parasite levels and body condition in the promiscuous Aquatic Warbler (*Acrocephalus paludicola*), 558–565

Eagle, Bald, 994–999; Crested Serpent, 108–120
eavesdropping, 94–107
ecological, – barriers, 1281–1290; – niche, 689–700; – plasticity, 689–700
ecomorphology, 689–700
economics of defense, 187–204
Ectopistes migratorius, 389–402, 1003–1004
Edelaar, Pim, et al., No support for a genetic basis of mandible crossing direction in crossbills (*Loxia* spp.), 1123–1129
edge, 618–636; – avoidance, 618–636; – effect, 36–49
Edwards, Scott V., et al., Future of avian genetic resources collections: Archives of evolutionary and environmental history, 979–984
effort-adjustment factors, 338–346
egg, – formation, 1239–1248; – investment, 1161–1168; – measurements, 994–999; – production, 313–318; – quality, 1239–1248; replacement –, 566–570; – size, 246–254, 509–522
eggshell, – collections, 994–999; – fragments, 994–999; – thickness, 994–999; – thinning, 990–994
Egret, Great, 255–267
Eider, Common, 523–529; King, 313–318
El Niño-Southern Oscillation, 571–592
Emberiza, 733–746
E. calandra, 853–871
E. hortluna, 733–746
E. rustica, 733–746
Emberizidae, 1–14, 733–746, 938–948, 1026–1031
Emberizinae, 1026–1031
embryo, – mortality, 1239–1248; – sex, 1239–1248
emigration, 205–221
Emlen, Stephen T., see Wrege, Peter H., and –
Empidonax, 1070–1085
E. alnorum, 544–557, 689–700, 1070–1085
E. difficilis, 938–948, 1070–1085
E. flaviventris, 938–948
E. hammondii, 689–700, 938–948, 1070–1085
E. minimus, 71–81, 689–700, 938–948, 1130–1148
E. oberholseri, 938–948, 1070–1085
E. occidentalis, 1070–1085
E. traillii, 1070–1085
E. t. extimus, 1261–1270
E. wrightii, 938–948, 1070–1085
endangered species, 1261–1270
endemism, 121–143, 1271–1280
endocrine, 255–267; – disruption, 915–921
endogenous reserves, 313–318, 313–318
enemy recognition, 530–543
energetics, 544–557; annual cycle –, 922–937
environmental, – monitoring, 966–971; – quality, 509–522
Eocene, 1037–1048
Epthianura albifrons, 280–291
Eremophila alpestris, 1130–1148
Errata, 378, 1023, 1309
Escamilla, Erica M., see Morris, Sara R., et al.
esophageal content, 144–157
estradiol, 82–93
Eubucco bourcierii, 1055–1063
Eucalyptus, 1097–1111
Eudocimus albus, 255–267
Eudynamys taitensis, 733–746
Eugenes fulgens, 1097–1111
Eulampis jugularis, 187–204
Euphagus cyanocephalus, 82–93
Euplectes, 1271–1280
E. orix, 57–70
Eupsittula, 292–305
Eurotrochilus inexpectatus, 1049–1054
Eurylaimidae, 1271–1280
Eurystomus gularis, 1271–1280
Evans, Roger Malcolm, 1935–1998, In Memoriam, 351–353
evolution, character –, 1191–1209; – cultural, 853–871; – molecular, 50–56
Ewert, David N., see Mehlman, David W., et al.
extrapair, – copulations, 57–70; – fertilization, 759–772, 1086–1096; – offspring, 747–758; – paternity, 57–70

Faaborg, John, review by, 373–375
Fairy-wren, Superb, 1086–1096, 1169–1181
Falco aeruginosus, 1026–1031
F. peregrinus, 994–999
F. sparverius, 71–81, 835–842, 1130–1148
F. tinnunculus, 1026–1031
Falcon, Peregrine, 994–999
Falconiformes, 971–979, 1130–1148, 1210–1224
fall migration, 319–328
family groups, 479–496
Farnsworth, Andrew, Flight calls and their value for future ornithological studies and conservation research, 733–746
Farnsworth, George L., and Simons, Theodore R., Relationship between Mayfield nest-survival estimates and seasonal fecundity: A cautionary reply, 1000–1001
fat, – deposition, 544–557; – loads, 922–937
feather-degrading bacteria, 222–230
feathers, 222–230, 255–267, 1037–1048; clipped –, 1225–1238
fecal contents, 783–792
fecundity, – models, 280–291, 306–312; seasonal –, 280–291, 306–312, 1000–1001, 1001–1003
feeding, 1249–1260; – ecology, 144–157; – rates, 1169–1181
female, – behavior, 57–70; – choice, 82–93; – quality, 902–914
fertility insurance, 57–70

Ficedula albicollis, 57–70, 509–522, 530–543, 558–565, 747–758, 829–834, 1112–1122
F. hypoleuca, 246–254, 509–522, 558–565, 689–700, 733–746, 747–758, 1130–1148, 1225–1238
field study, 819–828
Fieldfare, 733–746
Finch, 403–413, 733–746; Cassin's, 1070–1085; Darwin's, 747–758, 1210–1224; Galapagos, 403–413; House, 403–413; Purple, 1130–1148; Zebra, 1–14, 255–267, 509–522, 1112–1122, 1239–1248
Firecrown, Juan Fernández, 593–607
Fisher mechanism, 1–14
fitness, 1161–1168, 1239–1248
Fitzpatrick, John W., In Memoriam: Raymond Andrew Paynter, Jr., 1925–2003, 1295–1297
Fjeldså, Jon, The Grebes: Podicipedidae (rev.), 1301–1303
fledging success, 648–660, 1225–1238
Fleischer, Robert C., see Anderson, Kurt E., et al.
Flicker, 747–758; Northern, 1070–1085, 1130–1148
flight, – behavior, 872–886; – calls, 733–746; – performance, 872–886; – ranges, 922–937; – whistles, 82–93, 803–818
flight-calling behavior, 733–746
flowering, 1097–1111
Flycatcher, 733–746; 673–683, 938–948, 1070–1085; Alder, 544–557, 689–700, 1070–1085; Collared, 57–70, 509–522, 530–543, 558–565, 747–758, 829–834, 1112–1122; Cordilleran, 1070–1085; Dusky, 938–948, 1070–1085; Gray, 938–948, 1070–1085; Hammond's, 689–700, 938–948, 1070–1085; Least, 71–81, 689–700, 938–948, 1130–1148; Ochre-bellied, 783–792; Pacific-slope, 938–948, 1070–1085; Pied, 246–254, 509–522, 558–565, 689–700, 733–746, 747–758, 1130–1148, 1225–1238; Southwestern Willow, 1261–1270; Spotted, 733–746; Western, 1070–1085; Willow, 1070–1085; Yellow-bellied, 938–948
Foliage-gleaner, Henna-hooded, 389–402
food, – abundance, 783–792, 1225–1238; – allocation, 1112–1122; – availability, 144–157, 902–914; – competition, 1055–1063; – limitation, 1112–1122; – quality, 144–157; – resources, 313–318; – sample methods, 1182–1190; – supply, 1225–1238
food-hoarding capacity, 819–828
foraging, 1097–1111; – behavior, 36–49, 187–204; – ecology, 887–901; – efficiency, 1225–1238
Forbes, L. Scott, see Sealy, Spencer G., et al.
forest, 1281–1290; – fragmentation, 497–508
Forman, Dan W., Laying plasticity in an avian brood parasite, 566–570
Formicariidae, 673–683
Formicarius, 673–683
Forsyth, Mark H., see Cristol, Daniel A., et al.
Fort Union, 347–348
fossil, 1037–1048; – birds, 1055–1063; – specimens, 971–979
Foudia, 1271–1280
foudies, 1271–1280
fragmentation, 36–49, 618–636; forest –, 497–508
Francis, Charles M., see Dunn, Erica H., et al.
Fratercula arctica, 773–782, 887–901
Frederick, Peter C., see Heath, Julie A., and –
Fredrickson, Leigh H., and Havera, Stephen P., In Memoriam: Frank C. Bellrose, 1916–2005, 1298–1300
Friedl, Thomas W. P., and Klump, Georg M., Extrapair fertilizations in Red Bishops (*Euplectes orix*): Do females follow conditional extrapair strategies?, 57–70
fringillids, 733–746
Frith, Clifford B., and Frith, Dawn W., The Bowerbirds: Ptilonorhynchidae (rev.), 718–721
Frith, Dawn W., see Frith, Clifford B., and –
fruit biomass, 783–792
Fulica americana, 26–35, 566–570, 1130–1148
Fuller, Robert J., see Gillings, Simon, et al.
Fulmar, Atlantic Northern, 990–994
Fulmarus glacialis, 990–994
Funambulus, 108–120
Furnariidae, 158–174, 433–444, 673–683

Gadwall, 1130–1148
Galápagos Islands, 1210–1224
Galbulae, 1055–1063
Galliformes, 246–254, 971–979; 1049–1054
Gallinago delicata, 1130–1148
G. gallinago, 1149–1160
Gallinula chloropus, 26–35, 566–570
Gallus, 1161–1168
G. domesticus, 835–842, 843–852, 1239–1248
G. gallus, 121–143, 255–267, 949–965, 987–990, 1161–1168
G. g. domesticus, 158–174, 414–432
game bird, 1149–1160
Gammon, David E., et al., Cultural divergence within novel song in the Black-capped Chickadee (*Poecile atricapillus*), 853–871
Garrulax cinereifrons, 108–120
gastrointestinal tract, 887–901
Gaunt, Sandra L. L., et al., New directions for bioacoustics collections, 984–987
Gauthier, Gilles, see Cadieux, Marie-Christine, et al.
Gauthier, Gilles, see Menu, Stéphane, et al.
Gavia immer, 255–267
Gaviiformes, 971–979, 1055–1063
Gay, Laurène, et al., Additional data on mitochondrial DNA of North American large gull taxa, 684–688
gene flow, 329–337, 403–413, 803–818, 1149–1160, 1191–1209, 1210–1224
general improvement hypothesis, 509–522
general linear mixed model, 1161–1168
genetic, – basis, 1123–1129; – bottleneck, 464–478; – cline, 759–772; – differentiation, 464–478; – distance, 94–107; – divergence, 121–143;

– diversity, 94–107, 464–478, 1210–1224; – drift, 1210–1224; – index, 747–758, 759–772; – monomorphism, 1210–1224; – resources collections, 979–984; – structure, 949–965
genetics, population –, 464–478, 1210–1224
genotypes, 747–758, 759–772
Geobates, 158–174
G. poecilopterus, 158–174
Geococcyx californianus, 523–529
geographic, – structure, 949–965; – variation, 403–413, 648–660, 803–818, 853–871
Geositta, 158–174
G. antarctica, 158–174
G. crassirostris, 158–174
G. cunicularia, 158–174
G. c. cunicularia, 158–174
G. c. georgei, 158–174
G. excelsior, 158–174
G. isabellina, 158–174
G. maritima, 158–174
G. peruviana, 158–174
G. poecilopterus, 158–174
G. punensis, 158–174
G. rufipennis, 158–174
G. saxicolina, 158–174
G. tenuirostris, 158–174
Geospiza conirostris, 571–592
Geothlypis trichas, 1070–1085, 1130–1148
G. t. trichas, 793–802
Germany, 1055–1063
Geupel, Geoffrey R., see Chase, Mary K., et al.
Gibbs, H. Lisle, see Shriver, W. Gregory, et al.
Gibbs, James P., see Conway, Courtney J., and –
Gibbs, James P., see Shriver, W. Gregory, et al.
Gibson, Daniel D., see Kulikova, Irina V., et al.
Gilchrist, H. Grant, see DeVink, Jean-Michel A., et al.
Gill, Frank B., et al., Phylogeny of titmice (Paridae): II. Species relationships based on sequences of the mitochondrial cytochrome-*b* gene, 121–143
Gill, Sharon A., and Stutchbury, Bridget J. M., Nest building is an indicator of parental quality in the monogamous Neotropical Buff-breasted Wren (*Thryothorus leucotis*), 1169–1181
Gillings, Simon, et al., Diurnal studies do not predict nocturnal habitat choice and site selection of European Golden-Plovers (*Pluvialis apricaria*) and Northern Lapwings (*Vanellus vanellus*), 1249–1260
Given, Andrew D., et al., Molecular evidence for recent radiation in Southern Hemisphere masked gulls, 268–279
glacial events, 158–174
Glaucidium gnoma, 1070–1085
Glenn, Travis C., see Korfanta, Nicole M., et al.
Gloger's Rule, 222–230
Glossopsitta concinna, 1097–1111
Gnatcatcher, Blue-gray, 280–291; California, 571–592
gnateaters, 673–683
Godwit, Bar-tailed, 922–937; Marbled, 1130–1148
Goldeneye, Common, 313–318, 1130–1148
Golden-Plovers, European, 1249–1260
Goldenthroat, Tepui, 389–402
Goldman, Robin S., see Temeles, Ethan J., et al.
González, Clementina, and Ornelas, Juan Francisco, Song structure and microgeographic song variation in Wedge-tailed Sabrewings (*Campylopterus curvipennis*) in Veracruz, Mexico, 593–607
good genes, 57–70; – hypothesis, 509–522
good resources, 689–700
Goodale, Eben, and Kotagama, Sarath W., Alarm calling in Sri Lankan mixed-species bird flocks, 108–120
Goose, 144–157, 313–318, 479–496, 1037–1048; Barnacle, 479–496; Cackling, 144–157; Canada, 144–157, 523–529, 1037–1048, 1130–1148; Emperor, 479–496; Greater Snow, 144–157, 479–496; Greater White-fronted, 1130–1148; Hawaiian, 329–337; Lesser Snow, 144–157, 313–318, 479–496; Ross's, 313–318; Snow, 144–157, 1130–1148
Gorman, Helen E., et al., Effects of incubation conditions and offspring sex on embryonic development and survival in the Zebra Finch (*Taeniopygia guttata*), 1239–1248
Goshawk, Crested, 108–120
Goura, 1003–1004
Grackle, Common, 1130–1148
Grallaria, 673–683
G. alleni, 389–402
G. blakei, 673–683
G. dignissima, 673–683
G. eludens, 673–683
G. ruficapilla, 673–683
G. rufula, 673–683
G. squamigera, 673–683
G. varia, 673–683
G. watkinsi, 389–402, 673–683
Grallaricula, 673–683
G. flavirostris, 673–683
G. lineifrons, 673–683
G. peruviana, 389–402
Grallarinae, 673–683
Granfors, Diane A., see Pietz, Pamela J., and –
Grant, Todd A., et al., Time-specific variation in passerine nest survival: New insights into old questions, 661–672
grassland birds, 618–636
grazing, 144–157, 618–636
Great Plains, 648–660
Grebe, 15–25; Atitlan, 389–402; Horned, 1130–1148; Pied-billed, 26–35, 1130–1148; Red-necked, 1130–1148; Western, 1130–1148
Grim, Tomáš, Host recognition of brood parasites: Implications for methodology in studies of enemy recognition, 530–543

Grosbeak, 733–746; Black-headed, 1070–1085; Rose-breasted, 71–81, 280–291, 938–948
Ground-Jay, Tibetan, 121–143
group vigilance, 108–120
Grouse, Black, 71–81
growth, 523–529, 1003–1004
Grubb, Thomas C., Jr., see Bronson, C. L., et al.
Gruiformes, 971–979, 1130–1148
Grus canadensis, 1130–1148
Grzybowski, Joseph A., and Pease, Craig M., Renesting determines seasonal fecundity in songbirds: What do we know? What should we assume?, 280–291
Guarouba guarouba, 292–305
Gulf of Maine, 773–782
Gull, 15–25, 268–279, 684–688, 1182–1190; Andean, 268–279; Black-billed, 268–279; Black-headed, 268–279; Bonaparte's, 268–279; Brown-headed, 268–279; Brown-hooded, 268–279; Franklin's, 268–279, 1130–1148; Gray-hooded, 268–279; Great Black-backed, 915–921; Hartlaub's, 268–279; Herring, 255–267, 915–921, 1130–1148; Ivory, 15–25; Laughing, 268–279; Little, 268–279; Mediterranean, 268–279; Red-billed, 268–279; Ring-billed, 1130–1148; Saunders's, 268–279; Silver, 268–279; Slender-billed, 268–279
Gutscher, Hans, see Stokke, Bård G., et al.
Gwinner, Eberhard, 1938–2004, In Memoriam, 1013–1015
Gygnis alba, 1026–1031
Gymnopithys leucaspis, 843–852
Gymnopsittacus, 292–305
Gymnorhinus cyanocephalus, 36–49, 50–56

Habia fuscicauda,, 783–792
habitat, 497–508, 648–660, 1261–1270, 1281–1290; – characteristics, 497–508; – connectivity, 853–871; – quality, 902–914; – selection, 1097–1111, 1249–1260; – use, 144–157
Haematopus palliatus, 15–25
Haftorn, Svein, 1925–2003, In Memoriam, 358
Halcyon badia, 1271–1280
H. leucocephala, 1271–1280
H. malimbica, 1271–1280
H. senegalensis, 733–746
H. winchelli, 1271–1280
Haldane's rule, 759–772
Haliaeetus leucocephalus, 994–999
Hamao, Shoji, and Saito, Daichi S., Extrapair fertilization in the Black-browed Reed Warbler (*Acrocephalus bistrigiceps*): Effects on mating status and nesting cycle of cuckolded and cuckolder males, 1086–1096
handicap principle, 1–14
Hanner, Robert, see Edwards, Scott V., et al.
Hapalopsittaca fuertesi, 389–402
Hargitai, Rita, et al., Effects of environmental conditions and parental quality on inter- and intraclutch egg-size variation in the Collared Flycatcher (*Ficedula albicollis*), 509–522
Harpactes fasciatus, 108–120
Harper, David, Recognition of avian sibling species, 1291–1292
Harrier, Eastern Marsh, 1026–1031; Hen, 1291–1292; Montagu's, 1291–1292; Northern, 1130–1148; Western Marsh, 1026–1031
Harrigan, Ryan J., see Kulikova, Irina V., et al.
Hartlaubius auratus, 1271–1280
hatching, – failures, 57–70; – success, 246–254, 835–842, 1239–1248
Havera, Stephen P., see Fredrickson, Leigh H., and –
Hawk, 108–120; Cooper's, 1037–1048, 1130–1148; Galápagos, 1210–1224; Red-tailed, 1130–1148; Rough-legged, 1130–1148; Sharp-shinned, 1130–1148; Swainson's, 1210–1224
Heath, Julie A., and Frederick, Peter C., Relationships among mercury concentrations, hormones, and nesting effort of White Ibises (*Eudocimus albus*) in the Florida Everglades, 255–267
Hegyi, Gergely, see Hargitai, Rita, et al.
Heliconia, 187–204
helicopter dynamics model, 872–886
Helmitheros vermivorum, 280–291
Hemicircus concretus, 1055–1063
Henicorhina leucosticta, 50–56
heritability, 1123–1129
Hermit, Little, 593–607
Hernández C., Patricia, see Herrera M., L. Gerardo, et al.
Heron, Great Blue, 1130–1148; Green, 26–35
Herrera M., L. Gerardo, et al., Quantifying differential responses to fruit abundance by two rainforest birds using long-term isotopic monitoring, 783–792
heterophils, 829–834
high-latitude migrants, 544–557
hippocampal volume, 819–828
hippocampus, 608–617;
hirundines, 637–647
Hirundinidae, 1055–1063
Hirundo, 121–143
H. ariel, 1086–1096
H. rustica, 509–522, 637–647, 1130–1148
history of ornithology, 389–402
hoarding, food –, 819–828; food – intensity, 819–828
Hobson, Keith A., see Herrera M., L. Gerardo, et al.
Hobson, Keith A., Stable isotopes and the determination of avian migratory connectivity and seasonal interactions, 1048–1037
Hodgman, Thomas P., see Shriver, W. Gregory, et al.
Hoelzer, Guy A., see Arsenault, David P., et al.
Höfling, Elizabeth, see Silveira, Luís Fábio, et al.
Holberton, Rebecca L., and Able, Kenneth P., In Memoriam: Eberhard Gwinner, 1938–2004, 1013–1015
Holmes, Richard T., see Jones, Jason, et al.

Holmes, Richard T., see Nagy, Laura R., and –
home ranges, 36–49
homing, 608–617
honeycreepers, 187–204, 1210–1224
Honeyeater, 1097–1111; Brown-headed, 1097–1111; Black-chinned, 1097–1111; Fuscous, 1097–1111; White-eared, 1097–1111; White-naped, 1097–1111; Yellow-faced, 1097–1111; Yellow-plumed, 1097–1111; Yellow-tufted, 1097–1111
honeyguides, 1049–1054, 1055–1063
hormones, 255–267
Hornbill, African Grey, 1271–1280
Hosoi, S. Aki, et al., Sexual preferences of female Brown-headed Cowbirds (*Molothrus ater*) for perched song repertoires, 82–93
host attributes, 648–660
House-Martin, Common, 637–647
Houston, C. Stuart, 363–365
Howe, Marshall A., see Dunn, Erica H., et al.
Howell, Thomas Raymond, 1924–2004, In Memoriam, 1008–1010
hue, 793–802
Hughes, R. John, see Cadieux, Marie-Christine, et al.
Hummingbird, 187–204, 389–402, 445–463, 593–607, 853–871, 872–886, 938–948, 1049–1054, 1055–1063; Allen's, 872–886; Amethyst-throated, 593–607; Anna's, 187–204, 872–886; Black-chinned, 187–204, 872–886, 938–948, 1097–1111; Blue-throated, 593–607, 1097–1111; Blue-vented, 187–204; Calliope, 1070–1085; Rivoli's, 1097–1111; Ruby-throated, 872–886, 938–948; Rufous, 187–204, 872–886, 938–948, 1070–1085
Hunter, W. C., see Rich, T. D., et al.
hunting, 479–496
hybrid zone, 414–432, 433–444, 759–772
hybridization, 94–107, 268–279, 747–758, 949–965
Hylocharis grayi, 1026–1031
H. humboldtii, 1026–1031
Hylocichla mustelina, 280–291, 306–312
Hylocryptus erythrocephalus, 389–402
Hylopezus, 673–683
H. berlepschi, 673–683
H. fulviventris, 673–683
H. nattereri, 292–305, 673–683
H. ochroleucus, 292–305
H. persipicillatus, 843–852
Hylophylax naevioides, 445–463, 843–852
hyperphagia, 544–557
Hypopyrrhus pyrohypogaster, 445–463
Hypothymis azurea, 108–120
Hypsibemon, 673–683
Hypsipetes madagascariensis, 1271–1280

Ibis, 1049–1054; White, 255–267
Icteria virens, 938–948
icterid orioles, 938–948
Icterus bullockii, 1070–1085
I. cucullatus, 1070–1085
I. galbula, 938–948, 1070–1085, 1130–1148
I. laudabilis, 187–204
I. spurius, 938–948, 1070–1085
Idaho, 1070–1085
immigration, 205–221
immune, – function, 829–834; – system, 558–565
inbreeding, 464–478
income breeders, 1225–1238
incomplete lineage sorting, 949–965
incubation, 313–318, 835–842, 1239–1248
Indicator variegatus, 1055–1063
Indicatoridae, 1049–1054, 1055–1063
Iñigo-Elias, E. E., see Rich, T. D., et al.
insectivores, 246–254
insectivorous passerines, 246–254, 509–522
insemination, artificial –, 1161–1168
interbreed, 747–758
interclutch variation, 509–522
interdialect movement, 803–818
intraclutch variation, 509–522
introgression, 94–107, 414–432, 747–758, 759–772
invertebrates, 1261–1270
Iole indica, 108–120
Isler, Morton L., et al., Clinal variation in vocalizations of an antbird (Thamnophilidae) and implications for defining species limits, 433–444
Isler, Phyllis R., see Isler, Morton L., et al.
isoleg analysis, 1097–1111
isotopic landscape, 1037–1048
Ispidina, 1271–1280
I. lecontei, 1271–1280
I. madagascariensis, 1271–1280
I. picta, 1271–1280
Ixobrychus exilis, 26–35

Jabouilleia, 1064–1069
Jabouilleia danjoui, 1064–1069
J. d. danjoui, 1064–1069
J. d. parvirostris, 1064–1069
J. naungmungensis, 1064–1069
Jacana, Wattled, 347–348, 348–349
Jacana jacana, 329–337, 348–349
Jackson, Jerome A., review by, 716–718
Jacques, Christopher N., see Shriver, W. Gregory, et al.
James, Frances C., and Pourtless, John A., IV, review by, 714–716
James, Helen F., Paleogene fossils and the radiation of modern birds, 1049–1054
Jay, Gray, 36–49; Mexican, 571–592, 1130–1148; Pinyon, 36–49; Steller's, 36–49, 571–592, 1070–1085
Jehl, J. R., Jr., review by, 1020–1022
Jenkinson AOU Service Award, 2004: C. Stuart Houston, 363–365
Jensen, William E., and Cully, Jack F., Jr., Geographic variation in Brown-headed Cowbird (*Molothrus ater*) parasitism on Dickcissels (*Spiza americana*) in Great Plains tallgrass prairie, 648–660

Johnston, David W., The History of Ornithology in Virginia (rev.), 375–377
Jones, Jason, et al., Mayfield nest-survival estimates and seasonal fecundity: Reply to Farnsworth and Simons, 1001–1003
Jones, Jason, et al., Relationship between Mayfield nest-survival estimates and seasonal fecundity: A cautionary note, 306–312
Jones, Peter T., see Shriver, W. Gregory, et al.
Jones, Stephanie L., In Memoriam: George William Salt, 1919–1999, 354
Junco, Dark-eyed, 544–557, 608–617, 1070–1085, 1130–1148
Junco hyemalis, 544–557, 608–617, 1070–1085, 1130–1148
J. h. carolinensis, 608–617
J. h. hyemalis, 608–617
Junglefowl, Red, 987–990, 1161–1168
Jynginae, 1055–1063
Jynx ruficollis, 1055–1063
J. torquilla, 1055–1063

Kaltenecker, Gregory S., see Carlisle, Jay D., et al.
Kamil, Alan C., 359–361
Kear, Janet, 1933–2004, In Memoriam, 1293–1294
Keiser, Jeffrey T., et al., Homing success of migrant versus nonmigrant Dark-eyed Juncos (*Junco hyemalis*), 608–617
Keith, Allan R., et al., The Birds of Hispaniola (rev.), 1016–1018
Kellett, Dana K., et al., Body mass of Long-tailed Ducks (*Clangula hyemalis*) during incubation, 313–318
Kennedy, J. A., see Rich, T. D., et al.
keratin, 222–230
Kern, Michael D., see Owen, Jennifer C., et al.
Kestrel, American, 71–81, 835–842, 1130–1148
Kiff, Lloyd F., History, present status, and future prospects of avian eggshell collections in North America, 994–999
Kiff, Lloyd F., see Cade, Tom J., et al.
Killdeer, 1130–1148
Kingbird, Eastern, 1130–1148; Western, 1130–1148
Kingfisher, 1271–1280; African Dwarf, 1271–1280; African Malachite, 1271–1280; African Pygmy, 1271–1280; Belted, 1130–1148; Black-backed, 1271–1280; Blue-breasted, 1271–1280; Blue-eared, 1271–1280; Chocolate-backed, 1271–1280; Collared, 1271–1280; Common, 1271–1280; Dwarf, 1271–1280; Dwarf River, 1271–1280; Giant, 1271–1280; Green, 1271–1280; Grey-headed, 1271–1280; Madagascar Malachite, 1271–1280; Madagascar Pygmy, 1271–1280; Malachite, 1271–1280; Philippine, 1271–1280; Pied, 1271–1280; Rufous-backed, 1271–1280; Rufous-lored, 1271–1280; Shining Blue, 1271–1280; Variable, 1271–1280; White-bellied, 1271–1280; Woodland, 733–746
Kinglet, 938–948; Golden-crowned, 1070–1085; Ruby-crowned, 71–81, 544–557, 689–700, 938–948, 1070–1085, 1130–1148
Kittiwake, Black-legged, 268–279
Klump, Georg M., see Friedl, Thomas W. P., and –
Knops, Peter, see Edelaar, Pim, et al.
Knot, Great, 922–937; Red, 922–937, 1249–1260
Koel, Long-tailed, 733–746
Komar, Oliver, et al., Evidence of latitudinal sexual segregation among migratory birds wintering in Mexico, 938–948
Koppelhus, Eva B., see Currie, Phillip J., et al.
Korfanta, Nicole M., et al., Burrowing Owl (*Athene cunicularia*) population genetics: A comparison of North American forms and migratory habits, 464–478
Kotagama, Sarath W., see Goodale, Eben, and –
Krabbe, Niels, see Cuervo, Andrés M., et al.
Kratter, Andrew W., review by, 712–714
Kress, Stephen W., see Breton, André R., et al.
Kruszewicz, Andrzej, see Dyrcz, Andrzej, et al.
Kudla, Alexei U., see Temeles, Ethan J., et al.
Kulikova, Irina V., et al., Phylogeography of the Mallard (*Anas platyrhynchos*): Hybridization, dispersal, and lineage sorting contribute to complex geographic structure, 949–965

Lampornis amethystinus, 593–607
L. clemenciae, 593–607, 1097–1111
Lance, Monique M., and Thompson, Christopher W., Overlap in diets and foraging of Common Murres (*Uria aalge*) and Rhinoceros Auklets (*Cerorhinca monocerata*) after the breeding season, 887–901
landbird migrants, 1070–1085
landbirds, 15–25, 319–328, 1049–1054, 1070–1085, 1130–1148, 1281–1290
landscape, 497–508, 1281–1290
Lanius ludovicianus, 1037–1048
Lapwing, Northern, 1249–1260
Laridae, 268–279
Lark, 733–746; Horned, 1130–1148
Larracuente, Amanda M., see Morris, Sara R., et al.
Larsson, Hans, see Olsen, Klaus Malling, and –
Larus, 268–279, 684–688
L. argentatus, 255–267, 684–688, 915–921, 1130–1148
L. atricilla, 268–279
L. brunnicephalus, 268–279
L. bulleri, 268–279
L. californicus, 684–688
L. canus, 684–688
L. cirrocephalus, 268–279
L. c. cirrocephalus, 268–279
L. c. poiocephalus, 268–279
L. crassirostris, 684–688
L. delawarensis, 1130–1148
L. dominicanus, 684–688
L. fuscus, 684–688

L. genei, 268–279
L. glaucescens, 684–688
L. glaucoides, 684–688
L. g. kumlieni, 684–688
L. hartlaubii, 268–279
L. hyperboreus, 684–688
L. maculipennis, 268–279
L. marinus, 684–688, 915–921
L. melanocephalus, 268–279
L. michahellis, 684–688
L. minutus, 268–279
L. novaehollandiae, 268–279
L. n. hartlaubii, 268–279
L. n. novaehollandiae, 268–279
L. n. scopulinus, 268–279
L. occidentalis, 684–688
L. o. occidentalis, 684–688
L. o. wymani, 684–688
L. philadelphia, 268–279
L. pipixcan, 268–279, 1130–1148
L. ridibundus, 268–279
L. saundersi, 268–279
L. schistisagus, 684–688
L. serranus, 268–279
L. smithsonianus, 684–688
L. thayeri, 684–688
laser engraving, 971–979
Laterallus jamaicensis, 26–35
Latta, Steven C., see Keith, Allan R., et al.
Laughingthrush, Ashy-headed, 108–120
Lawson, Shona L., see Kellett, Dana K., et al.
laying date, 1225–1238; – order, 509–522; – patterns, 566–570; – plasticity, 566–570
Leader, Noam, et al., Acoustic properties of two urban song dialects in the Orange-tufted Sunbird (*Nectarinia osea*), 231–145
Leisler, Bernd, see Dyrcz, Andrzej, et al.
leks, 593–607
Lepage, Denis, see Dunn, Erica H., et al.
Leptotila battyi, 1026–1031
L. plumbeiceps, 1026–1031
L. rufaxilla, 1026–1031
Lichenostomus, 1097–1111
L. chrysops, 1097–1111
L. fuscus, 1097–1111
L. leucotis, 1097–1111
L. melanops, 1097–1111
L. ornatus, 1097–1111
Liebner, David A., see Morris, Sara R., et al.
life history, 661–672
light conditions, 1249–1260
Limnothlypsis swainsonii, 689–700
Limosa fedoa, 1130–1148
L. lapponica baueri, 922–937
L. l. taymyrensis, 922–937
Limpkin, 26–35
Liosceles, 673–683
L. thoracicus, 673–683
Lipaugus weberi, 445–463
Locustella, 706–707
L. fluviatilis, 733–746
L. lusciniodes, 1291–1292
L. naevia, 1291–1292
logistic exposure, 661–672
Lonchura, 1271–1280
Long Point Bird Observatory, 71–81
Longspur, Chestnut-collared, 701–705
long-term, – study, 571–592, 783–792, 1130–1148; – trends, 15–25
Loon, 15–25, 1055–1063; Common, 255–267
Lophodytes cucullatus, 1130–1148
Lophophanes, 121–143
L. cristatus, 121–143
L. dichrous, 121–143
Lorikeet, Musk, 1097–1111
loudsong pace, 433–444
Loxia, 1123–1129, 1225–1238
L. curvirostra, 1123–1129
L. c. curvirostra, 1123–1129
L. c. himalayensis, 1123–1129
L. leucoptera bifasciata, 1123–1129
L. leucoptera leucoptera, 1123–1129
L. pytyopsittacus, 1123–1129
L. scotica, 1123–1129
Loxigilla noctis, 187–204
Luscinia svecica, 71–81
Lybius minor, 1055–1063
L. vieilloti, 1055–1063
Lyrurus tetrix, 71–81

Mabey, Sarah E., see Mehlman, David W., et al.
Mac Nally, Ralph, and Timewell, Christopher A. R., Resource availability controls bird-assemblage composition through interspecific aggression, 1097–1111
Machlolophus, 121–143
Madagascar, 1271–1280
Madden, Elizabeth M., see Grant, Todd A., et al.
Magpie-Robin, Madagascar, 1271–1280
male quality, 1–14, 57–70
Malkoha, Red-faced, 108–120
Mallard, 255–267, 509–522, 523–529, 558–565, 949–965, 1130–1148
Malurus cyaneus, 1086–1096, 1169–1181
Manacus manacus, 403–413
management, 15–25, 1149–1160
Manakin, White-bearded, 403–413
mandible crossing direction, 1123–1129
Manitoba, 1130–1148
Marchetti, Karen, see McNett, Gabriel D. and –
mark–recapture, 329–337, 803–818
Marks, Ben D., and Willard, David E., Phylogenetic relationships of the Madagascar Pygmy Kingfisher (*Ispidina madagascariensis*), 1271–1280
Marks, Jeffrey S., and Nordhagen, Ted, Type locality of *Ammodramus bairdii* (Audubon), 349–350

marsh birds, 15–25, 26–35
Martell, A. M., see Rich, T. D., et al.
Martin, Crag, 121–143; Fairy, 1086–1096; House, 121–143; Purple, 558–565, 1130–1148
Martínez Gómez, Juan E., et al., Phylogenetic position and generic placement of the Socorro Wren (*Thryomanes sissonii*), 50–56
Marzluff, John M., see Vigallon, Stacey M., and –
Marzluff, John M., see Withey, John C., and –
mass distributions, 347–348
mate, – choice, 1–14, 71–81, 82–93, 1169–1181; – guarding, 1086–1096
maternal effects, 509–522
mating, – status, 1086–1096; – success, 82–93; – system, 57–70, 558–565, 1210–1224
maximum size, 1003–1004
Mayfield nest-survival estimates, 306–312, 1000–1001, 1001–1003
Mayfield's estimator, 661–672
Mayr, Ernst, 1904–2005, In Memoriam, 1005–1007
Mayr, Gerald, A tiny barbet-like bird from the Lower Oligocene of Germany: The smallest species and earliest substantial fossil record of the Pici (woodpeckers and allies), 1055–1063
McAuley, Daniel G., see Rhymer, Judith M., et al.
McCracken, Kevin G., see Kulikova, Irina V., et al.
McDonald, David B., see Korfanta, Nicole M., et al.
McGillivray, W. Bruce, In Memoriam: Philip H. R. Stepney, 1947–2004, 711
McNett, Gabriel D., and Marchetti, Karen, Ultraviolet degradation in carotenoid patches: Live versus museum specimens of wood warblers (Parulidae), 793–802
Meadowlark, 618–636; Eastern, 618–636, 648–660, 661–672; Western, 82–93, 661–672, 1130–1148
Megaceryle alcyon, 1271–1280
M. maxima, 1271–1280
Megalaima, 1055–1063
M. armillaris, 1055–1063
M. asiatica, 1055–1063
M. javensis, 1055–1063
M. mystacophanos, 1055–1063
M. virens, 1055–1063
Megarhynchus pitangua, 1055–1063
Mehl, Katherine R., see Kellett, Dana K., et al.
Mehlman, David W., et al., Conserving stopover sites for forest-dwelling migratory landbirds, 1281–1290
Melanerpes carolinus, 558–565, 1055–1063
M. chrysauchen, 1026–1031
M. pulcher, 1026–1031
Melaniparus, 121–143
Melanochlora, 121–143
M. sultanea, 121–143
M. s. gayeti, 121–143
M. s. sultanea, 121–143
Meleagris gallopavo, 618–636
Melithreptus, 1097–1111
M. brevirostris, 1097–1111
M. gularis, 1097–1111
M. lunatus, 1097–1111
Melopsittacus undulates, 733–746
Melospiza georgiana, 1–14
M. lincolnii, 544–557, 938–948, 1070–1085
M. melodia, 1–14, 82–93, 280–291, 403–413, 571–592, 648–660, 853–871, 1130–1148
M. m. gouldii, 571–592
Melphagidae, 1097–1111
Mennill, Daniel J., and Vehrencamp, Sandra L., Sex differences in singing and duetting behavior of Neotropical Rufous-and-white Wrens (*Thryothorus rufalbus*), 175–186
Menu, Stéphane, et al., Survival of young Greater Snow Geese (*Chen caerulescens atlantica*) during fall migration, 479–496
Menura novaehollandiae, 1055–1063
Menuridae, 1055–1063
Merganser, Common, 1130–1148; Hooded, 1130–1148; Red-breasted, 1130–1148
Mergus merganser, 1130–1148
M. serrator, 1130–1148
Meropidae, 733–746
Merops bulocki, 1271–1280
M. pusillus, 1271–1280
M. superciliosus, 1271–1280
meta-analysis, 26–35
methodology, 530–543
methylmercury, 255–267
microbes, 222–230
microhabitat, 497–508
microparasites, 558–565
microsatellite, 94–107; – DNA markers, 464–478
migrants, 608–617; long-distance –, 71–81, 1130–1148; Nearctic-Neotropical –, 319–328, 544–557; Neotropical –, 329–337; short-distance –, 71–81, 1130–1148; western –, 1070–1085
migration, 319–328, 464–478, 479–496, 544–557, 608–617, 637–647, 733–746, 922–937, 1130–1148, 1281–1290; autumn – timing, 1070–1085; differential –, 71–81, 938–948; fall –, 319–328; – flyways, 1149–1160; Nearctic-Neotropical –, 938–948; nocturnal –, 733–746 spring – arrival dates, 1130–1148
migratory, – birds, 71–81, 1037–1048; – connectivity, 1037–1048
Mills, Alexander M., Can breeding habitat be sexually selected?, 689–700
Mills, Alexander M., Protogyny in autumn migration: Do male birds "play chicken"?, 71–81
Mills, James A., see Given, Andrew D., et al.
mimicry, 108–120
Mimidae, 733–746, 1026–1031
Mimodes, 1026–1031
M. graysoni, 1026–1031
Mimus, 1026–1031
M. gilvus, 1026–1031

M. graysoni, 1026–1031
M. polyglottos, 71–81, 1026–1031
mining, 497–508
minisatellite DNA markers, 1210–1224
Miocene, 1037–1048
Mionectes oleagineus, 783–792
Mniotilta varia, 793–802, 938–948
Mirafra, 1271–1280
mitochondrial, – DNA, 268–269, 684–688, 1149–1160, 1191–1209, 1271–1280; – DNA sequence, 268–279, 445–463; – variation, 414–432
mixed-species flocks, 108–120, 1249–1260
mobbing, 108–120
Mock, D. W., More Than Kin and Less Than Kind: The Evolution of Family Conflict (rev.), 366–367
Mock, Douglas W., see Bartlett, Terri L., et al.
Mockingbird, Northern, 71–81; Socorro, 1026–1031
molecular, – characters, 158–174, 673–683; – evolution, 50–56; – markers, 747–758; – phylogenetics, 979–984
Møller, Anders Pape, see Stokke, Bård G., et al.
Molothrus ater, 82–93, 530–543, 571–592, 618–636, 648–660, 701–705, 803–818, 1130–1148
M. a. artemisae, 803–818
M. a. ater, 803–818
M. a. obscurus, 82–93, 803–818
molt, 222–230; – constraint, 71–81; – strategy, 1070–1085; flight-feather –, 1070–1085; prebasic –, 1070–1085; preformative –, 1070–1085
Monarch, Black-naped, 108–120
monitoring, 15–25, 26–35; – program, 338–346
monogamous males, 1086–1096
Monticola, 1271–1280
Moorhen, Common, 26–35, 566–570
morphological, 94–107; – characters, 268–279, 347–348, 673–683; – stasis, 121–143; – traits, 1123–1129
morphology, 121–143, 747–758, 971–979, 1037–1048, 1055–1063, 1191–1209; bill –, 187–204; sternal –, 673–683; wing –, 872–886
Morris, Ralph D., see Sealy, Spencer G., et al.
Morris, Sara R., et al., Multiple-day constancy as an alternative to pooling for estimating mark–recapture stopover length in Nearctic–Neotropical migrant landbirds, 319–328
mortality, 479–496, 523–529; embryo –, 1239–1248
Motacilla alba, 1026–1031
M. alba leucopsis, 1026–1031
M. a. ocularis, 1026–1031
M. capensis, 1271–1280
M. clara, 1271–1280
M. flaviventris, 1271–1280
M. lugens, 1026–1031
M. trochilus, 706–707
Motacillidae, 733–746
Motis, Anna, see Delacour, Jean, et al.
mousebirds, 1049–1054, 1055–1063
multiple, – brooding, 902–914; – nesting attempts, 1000–1001
Mumme, Ronald L., review by, 366–367
Murphy-Klassen, Heather M., et al., Long-term trends in spring arrival dates of migrant birds at Delta Marsh, Manitoba, in relation to climate change, 1130–1148
Murray, Peter F., and Vickers-Rich, Patricia, Magnificent Mihirungs. The Colossal Flightless Birds of the Australian Dreamtime (rev.), 367–371
Murre, Common, 773–782, 887–901; Thick-billed, 887–901
Murrelet, Marbled, 36–49, 990–994
Muscicapa striata, 733–746
Muscicapidae, 733–746
museum, 389–402, 793–802; – collections, 938–948, 971–979, 987–990; – specimens, 966–971, 990–994
Myadestes townsendi, 1070–1085
Myanmar, 1064–1069
Myiocyx, 1271–1280
M. lecontei, 1271–1280
Myna, Crested, 1026–1031
Myornis senilis, 445–463
Myriachus, 673–683
Myrmornis, 673–683
M. torquata, 673–683
Myrmothera, 673–683
M. campanisona, 673–683
M. simplex, 673–683
Myrmotherinae, 673–683
Myrmotherula fulviventris, 843–852

Nack, Jamie L., see Renfrew, Rosalind B., et al.
Nager, Ruedi G., see Gorman, Helen E., et al.
Nagy, Laura R., and Holmes, Richard T., To double-brood or not? Individual variation in the reproductive effort in Black-throated Blue Warblers (*Dendroica caerulescens*), 902–914
Nagy, Laura R., see Jones, Jason, et al.
Nandayus nenday, 292–305
Nannus, 50–56
N. troglodytes, 50–56
Natunaornis gigura, 1003–1004
natural selection, frequency-dependent –, 1123–1129
Navarro-Sigüenza, Adolfo G., see Komar, Oliver, et al.
navigation, 608–617
Nearctic species, 733–746
Nearctic–Neotropical migrants, 1281–1290
nectar, 1097–1111
Nectarinia, 1271–1280
N. osea, 231–245
nectarivore, 872–886, 1097–1111
nectarivorous birds, 187–204
Nelson, Douglas A., see Gaunt, Sandra L. L., et al.
Neornithes, 1049–1054
Neornithine, 1049–1054
Neotropical, – birds, 445–463; – migrants, 15–25, 71–81, 1070–1085; – songbirds, 36–49; – species, 733–746

Neotropics, 938–948
Nesillas, 1271–1280
nest, – attendance, 255–267; – building, 1169–1181; – defense, 530–543; – density, 618–636; –initiation, 661–672; – predation, 36–49, 280–291, 306–312, 571–592, 618–636, 661–672, 843–852; – productivity, 280–291; – success, 280–291, 571–592, 1001–1003; – survival, 618–636, 661–672
nesting, – cycle, 1086–1096; – effort, 255–267
nestling, 1112–1122; – position, 1112–1122; – provisioning, 1169–1181; size-monomorphic –, 1112–1122
nests, artificial –, 843–852; dormitory –, 1169–1181
neuroanatomy, 175–186
neuroecological approach, 819–828
New Mexico, 329–337
new species, 445–463, 1064–1069
niche, – partitioning, 887–901; – specialization, 158–174
Nicholson, Edward Max, 1904–2003, In Memoriam, 357
Nighthawk, Common, 1130–1148
Night-Heron, Black-crowned, 1130–1148
Nisbet, Ian C. T., Criteria for sexing birds in studies of sexual dimorphism and sex-specific behavior, 347–348
nitrogen-15, 783–792
nocturnal, 733–746, 1249–1260
noise, anthropogenic –, 231–245
nonbreeding season, 938–948
nonmigrants, 608–617
nonpasserine, 1049–1054
Nooker, Jacqueline K., et al., Effects of food abundance, weather, and female condition on reproduction in Tree Swallows (*Tachycineta bicolor*), 1225–1238
Nordhagen, Ted, see Marks, Jeffrey S., and –
North American Coordinated Bird Monitoring Plan, 15–25
North Atlantic Oscillation, 637–647
Nowicki, Stephen, and Searcy, William A., Song and mate choice in birds: How the development of behavior helps us understand function, 1–14
Nuechterlein, Gary L., review by, 1301–1303
Numenius americanus, 15–25
Numida meleagris, 1026–1031
Nunavik, 144–157
Nur, Nadav, see Chase, Mary K., et al.
Nuthatch, Red-breasted, 1070–1085
nutrient limitation, 246–254
nutritional, – ecology, 783–792, 1037–1048; – stress hypothesis, 1–14
Nycticorax nycticorax, 1130–1148

O'Loghlen, Adrian L., see Anderson, Kurt E., et al.
O'Loghlen, Adrian L., see Hosoi, S. Aki, et al.
O'Shea, B. J., see Komar, Oliver, et al.
observer bias, 26–35
Oceanodroma monorhis, 733–746
Oena capensis, 1271–1280
offspring quality, 509–522
Ohio, 747–758, 759–772
Oligocene, 1037–1048, 1055–1063
Olsen, Klaus Malling, and Larsson, Hans, Gulls of North America, Europe, and Asia (rev.), 1020–1022
Olson, Storrs L., review by, 367–371
oology, 994–999
open population models, 319–328
Oporornis formosus, 938–948, 1037–1048
O. tolmiei, 938–948, 1070–1085
Oriole, 747–758; Baltimore, 938–948, 1070–1085; 1130–1148; Bullock's, 1070–1085; Hooded, 1070–1085; St. Lucia, 187–204; Orchard, 938–948, 1070–1085
ornamentation, paternal –, 1161–1168
Ornelas, Juan Francisco, see González, Clementina, and –
ORNIS, 987–990
Oropendola, Chestnut-headed, 389–402
Oropezus, 673–683
Orr, H. Allen, see Coyne, Jerry A., and –
Orr, Kate J., see Gorman, Helen E., et al.
oscines, 1–14
osteology, 971–979
Ottenwalder, José A., see Keith, Allan. R., et al.
Otus, 1271–1280
O. flammeolus, 329–337
Ovenbird, 280–291, 793–802, 938–948
overdispersion, 773–782
Overview, 15–25, 403–413, 747–758, 1049–1054
Owen, Jennifer C., et al., Habitat and sex differences in physiological condition of breeding Southwestern Willow Flycatchers (*Empidonax traillii extimus*), 1261–1270
Owl, 558–565, 1049–1054; Barn, 733–746; Burrowing, 464–478; Flammulated, 329–337; Marsh, 1271–1280; Northern Spotted, 329–337; Short-eared, 1130–1148
Oxyura jamaicensis, 1130–1148
Oystercatcher, American, 15–25

Pagophila eburnea, 15–25
Palearctic species, 733–746
Paleogene, 1037–1048
Paleotropical species, 733–746
Panama, 843–852
Panjabi, A. O., see Rich, T. D., et al.
Panurus biarmicus, 689–700
Parakeet, 292–305; Canary-winged, 292–305; Carolina, 389–402; Golden, 292–305; Golden-capped, 292–305; Jandaya, 292–305; Nanday, 291–305; Peach-fronted, 292–305; **Sulfur-breasted**, 292–305; Sun, 292–305; White-necked, 389–402
paraphyly, 949–965
parasites, blood –, 558–565; micro-, 558–565

parasitism, brood –, 280–291, 530–543, 648–660; conspecific brood –, 566–570
Pardaliparus, 121–143
P. amabilis, 121–143
P. venustulus, 121–143
parental, – behavior, 1239–1248; – care, 835–842, 1112–1122, 1169–1181; – nest defense, 701–705; – quality, 1169–1181
Paridae, 121–143, 747–758, 819–828, 1023
Parinae, 121–143
Parker, Patricia G., see Bollmer, Jennifer L., et al.
Parker, Timothy H., No evidence for adaptive differential sex allocation in Red Junglefowl (*Gallus gallus*), 1161–1168
Parrot, 292–305, 853–871, 1049–1054, 1097–1111; Indigo-winged, 389–402
Parulidae, 71–81, 793–802,938–948
Parus, 121–143, 747–758, 1225–1238
P. afer, 121–143
P. a. thruppi, 121–143
P. albiventris, 121–143
P. amabilis, 121–143
P. ater, 121–143
P. a. aemodius, 121–143
P. a. ater, 121–143
P. a.-melanolophus clade, 121–143
P. atricapillus, 121–143
P. a. atricapillus, 121–143
P. atricristatus, 121–143
P. bicolor, 121–143
P. b. bicolor, 121–143
P. bokharensis, 747–758
P. caeruleus, 57–70, 121–143, 509–522, 558–565, 793–802, 1086–1096
P. c. caeruleus, 121–143
Parus carolinensis, 121–143
P. c. carolinensis, 121–143
P. c. extimus, 121–143
P. cinctus, 121–143, 819–828
P. c. cinctus, 121–143
P. cristatus, 121–143
P. c. cristatus, 121–143
P. cyanus, 121–143
P. c. tianschanicus, 121–143
P. davidi, 121–143
P. dichrous, 121–143
P. d. dichroides, 121–143
P. elegans, 121–143
P. e. elegans, 121–143
P. e. mindanensis, 121–143
P. fasciiventer, 121–143
P. flavipectus, 121–143
P. funereus, 121–143
P. gambeli, 121–143
P. g. baileyi, 121–143
P. g. gambeli, 121–143
P. griseus, 121–143
P. holsti, 121–143
P. hudsonicus, 121–143
P. h. littoralis, 121–143
P. inornatus, 121–143
P. i. transpositus, 121–143
P. lugubris, 121–143
P. l. lugens, 121–143
P. major, 57–70, 71–81, 121–143, 231–245, 246–254, 558–565, 747–758, 793–802, 829–834, 902–914, 1130–1148
P. m. afrodite, 121–143
P. melanolophus, 121–143
P. montanus, 108–120, 121–143, 819–828
P. m. borealis, 121–143
P. m. songarus, 121–143
P. monticolus, 121–143
P. m. yunnanensis, 121–143
P. niger, 121–143
P. n. niger, 121–143
P. palustris, 121–143
P. p. brevirostris, 121–143
P. p. palustris, 121–143
P. ridgwayi, 121–143
P. rubidiventris, 121–143
P. rufescens, 121–143
P. r. rufescens, 121–143
P. rufiventris, 121–143
P. r. pallidiventris, 121–143
P. rufonuchalis, 121–143
P. sclateri, 121–143
P. s. rayi, 121–143
P. spilonotus, 121–143
P. superciliosus, 121–143
P. varius, 121–143
P. v. castaneoventris, 121–143
P. v. varius, 121–143
P. wollweberi, 121–143
P. w. phillipsi, 121–143
P. xanthogenys, 121–143
Pashley, D. N., see Rich, T. D., et al.
Passer domesticus, 57–70, 835–842, 1086–1096
Passerculus rostratus, 403–413
P. sandwichensis, 403–413, 544–557, 618–636, 701–705
P. s. beldingi, 403–413
P. s. princeps, 403–413
P. s. rostratus, 403–413
P. s. sanctorum, 403–413
Passerella, 403–413
P. iliaca, 530–543, 544–557, 1070–1085, 1130–1148
Passeriformes, 57–70, 673–683, 1049–1054, 1055–1063, 1130–1148
Passerina amoena, 938–948, 1070–1085
P. ciris, 938–948, 1070–1085
P. cyanea, 280–291, 733–746, 938–948, 1086–1096
passerine, 71–81, 246–254, 329–337, 403–413, 544–557, 558–565, 608–617, 618–636, 637–647, 661–672, 673–683, 689–700, 701–705, 733–746, 783–792, 819–828, 902–914, 994–999, 1037–1048, 1049–1054, 1055–1063, 1070–1085,

1086–1096, 1130–1148, 1112–1122, 1225–1238, 1271–1280
pasture, 618–636
Patagonia, 158–174
paternity, 1086–1096
Pavo cristatus, 971–979
Paynter, Raymond Andrew, Jr., 1925–2003, In Memoriam, 1295–1297
Peafowl, 971–979
Pease, Craig M., see Grzybowski, Joseph A., and –
Pelecaniformes, 971–979, 1130–1148
Pelecanus erythrorhynchos, 1130–1148
P. occidentalis, 994–999
Pelican, 389–402; American White, 1130–1148; Brown, 994–999
Penguin, 1037–1048, 1049–1054; Adélie, 523–529, 1149–1160
perched songs, 82–93
perching birds, 1049–1054
Periparus, 121–143
P. amabilis, 121–143
P. ater, 747–758
P. a. aemodius, 121–143
P. a. ater, 121–143
P. elegans elegans, 121–143
P. e. mindanensis, 121–143
P. melanolophus, 121–143, 747–758
P. rubidiventris, 121–143
P. rufonuchalis, 121–143
Perisoreus canadensis, 36–49
Perrins, Christopher M., In Memoriam: Edward Max Nicholson, 1904–2003, 357
Perspectives in Ornithology, 1–14, 389–402, 733–746, 1037–1048
Petersen, Wayne R., and Meservey, W. Roger, Massachusetts Breeding Bird Atlas (rev.), 1305–1307
Peterson, A. Townsend, et al., Free and open access to bird specimen data: Why?, 987–990
Peterson, A. Townsend, see Komar, Oliver, et al.
Peterson, A. Townsend, see Martínez Gómez, Juan E., et al.
Petrochelidon, 121–143
P. pyrrhonata, 637–647, 1225–1238
Petronia petronia, 1086–1096
Pezophaps solitaria, 1003–1004
Phaenicophaeinae, 1271–1280
Phaenicophaeus pyrrhocephalus, 108–120
Phaethornis longuemareus, 593–607
Phalacrocorax aristotelis, 773–782
P. auritus, 1130–1148
Phalarope, Wilson's, 1130–1148
Phalaropus tricolor, 1130–1148
Phasianus colchicus, 618–636
Pheasant, Ring-necked, 618–636
phenotypic flexibility, 922–937
Pheucticus ludovicianus, 71–81, 280–291, 938–948
P. melanocephalus, 1070–1085
Pheugopedius fasciatoventris albigularis, 389–402
Philepittidae, 1271–1280
Phillips, Ron, see Edelaar, Pim, et al.
philopatry, 464–478
Phlegopsis, 673–683
P. nigromaculata, 673–683
Phoebe, Eastern, 1130–1148
Phoebetria fusca, 773–782
P. palpebrata, 773–782
Phoenicurus ochruros, 71–81
Phylloscartes chapmani, 389–402
Phylloscopus, 706–707, 1291–1292
P. collybita, 706–707, 733–746, 1291–1292
P. fuscatus, 71–81
P. sibilatrix, 1291–1292
P. trochilus, 530–543, 706–707, 1291–1292
phylogenetic, – analysis, 403–413; – history, 121–143; – position, 50–56; – relationships, 1271–1280; – species concept, 50–56, 403–413, 433–444
phylogenetics, 268–269; molecular –, 979–984
phylogeny, 121–143, 673–683, 684–688, 1191–1209, 1271–1280
phylogeography, 949–965, 979–984, 1149–1160
physiological condition, 1261–1270
physiology, 1261–1270; reproductive –, 255–267
Pici, 1049–1054, 1055–1063
Picidae, 1049–1054, 1055–1063
Piciformes, 971–979, 1049–1054, 1055–1063, 1130–1148, 1191–1209
Picinae, 1055–1063
Picoidea, 1055–1063
Picoides nuttallii, 1055–1063
P. pubescens, 108–120
piculets, 1055–1063
Picumninae, 1055–1063
Picumnus, 1055–1063
P. exilis, 1055–1063
Picus canus, 1055–1063
P. viridis, 1055–1063
Piersma, Theunis, see Battley, Phil F., and –
Pietz, Pamela J., and Granfors, Diane A., Parental nest defense on videotape: More reality than "myth", 701–705
Pietz, Pamela J., see Grant, Todd A., et al.
Pigeon, 608–617, 1003–1004; Nicobar, 1003–1004; Passenger, 389–402, 1003–1004; Rock, 530–543, 733–746
Pintail, Northern, 1130–1148
Pipilo, 403–413
P. maculatus, 1070–1085
Pipit, 733–746; Meadow, 530–543; Sprague's, 349–350
Piranga ludoviciana, 938–948, 1070–1085
P. rubra, 938–948
piscivorous birds, 255–267
Pitta, 733–746; Blue-winged, 733–746; Fairy, 733–746
Pitta moluccensis, 733–746
P. nympha, 733–746
Pittasoma, 673–683

P. michleri, 673–683
P. rufopileatum, 673–683
playback, 82–93, 231–245
Pleistocene, 158–174
Ploceidae, 57–70
Ploceus, 1271–1280
Plover, Mountain, 1037–1048; Piping, 15–25; Wilson's, 1249–1260
plumage, 158–174, 222–230, 347–348, 445–463, 673–683, 793–802, 915–921, 922–937, 1191–1209, 1271–1280; – characteristics, 94–107; – characters, 292–305, 403–413
Pluvialis apriacaria, 1249–1260
Podiceps auritus, 1130–1148
P. grisegena, 1130–1148
Podicipidiformes, 971–979, 1130–1148
Podilymbus gigas, 389–402
P. podiceps, 26–35, 1130–1148
Poecile, 121–143, 747–758
P. atricapillus, 108–120, 121–143, 689–700, 747–758, 759–772, 819–828, 853–871, 1070–1085
P. carolinensis, 747–758, 759–772
P. c. carolinensis, 121–143
P. c. extimus, 121–143
P. cincta, 747–758
P. cinctus, 121–143
P. davidi, 121–143
P. gambeli, 747–758, 853–871, 1070–1085
P. g. baileyi, 121–143
P. g. gambeli, 121–143
P. hudsonica, 121–143, 747–758
P. lugubris, 121–143
P. montanus, 121–143, 747–758
P. m. borealis, 121–143
P. m. songarus, 121–143
P. palustris, 747–758
P. p. brevirostris, 121–143
P. p. palustris, 121–143
P. rufescens, 82–93, 121–143
P. sclateri, 121–143, 747–758
P. superciliosus, 121–143
P. varius, 121–143
Pogoniulus, 1055–1063
P. bilineatus, 1055–1063
P. scolopaceus, 1055–1063
point counts, 15–25
point-count surveys, 26–35
Polemond River, 144–157
Polioptila caerulea, 280–291
P. californica, 571–592
polyandry, 1210–1224; cooperative –, 1210–1224
polychlorinated biphenyls, 915–921
polygynous males, 1086–1096
polygyny, 57–70
polymorphism, 1123–1129
Polytmus milleri, 389–402
Pomatorhinus, 1064–1069
P. gravivox, 1064–1069
P. tickelli, 1064–1069
ponderosa pine, 329–337
Pooecetes gramineus, 661–672, 701–705, 1070–1085, 1130–1148
population, – density, 571–592; – differentiation, 949–965; – dynamics, 205–221, 571–592, 661–672, 902–914; – expansion, 1149–1160; – genetics, 464–478, 1210–1224; island-, 1210–1224; – monitoring, 306–312; – size, 637–647; – structure, 329–337, 1149–1160; – trends, 338–346
Porzana carolina, 26–35, 1130–1148
Postma, Erik, see Edelaar, Pim, et al.
Pourtless, John A., IV, see James, Frances C. and –
precocial young, 523–529
predation, 661–672, 701–705; daily – rate, 843–852; incidental –, 36–49; nest –, 36–49, 280–291, 306–312, 571–592, 618–636, 661–672, 843–852
predator, 530–543, 618–636, 843–852; aerial –, 108–120, – vigilance, 108–120
premigratory hypertrophy, 922–937
prey, 1182–1190; – abundance, 887–901
probability of fledging, 1000–1001
Procellariformes, 971–979, 1055–1063
Progne subis, 558–565, 1130–1148
program MARK, 773–782
promiscuity, 558–565
protandry, 71–81
protein, – requirements, 783–792; – supplement, 1239–1248
protogyny, 71–81
Prunella modularis, 57–70
Psarocolius wagleri, 389–402
Pseudocalyptomena, 1271–1280
Pseudopodoces, 121–143
P. humilis, 121–143
Psilopogon, 1055–1063
P. pyrolophus, 1055–1063
Psittacara, 292–305
Psittacidae, 292–305
Psittaciformes, 292–305, 971–979, 1049–1054
Pteroglossus, 1055–1063
P. aracari, 1055–1063
P. castanotis, 1055–1063
P. inscriptus, 1191–1209
P. viridis, 1055–1063
Pteroptochos tarnii, 445–463
Ptyonoprogne, 121–143
Puffin, Atlantic, 773–782, 887–901
Pygmornis hermits, 872–886
Pygmy-owl, Northern, 1070–1085
Pygoscelis adeliae, 523–529, 1149–1160
Pyrenestes, 1123–1129
P. ostrinus, 1123–1129
Pyrrhura albipectus, 389–402

Quail, 843–852; Japanese, 36–49, 1037–1048
quality, egg –, 1239–1248; environmental –, 509–522; female –, 902–914; food –, 144–157; habitat –,

902–914; male –, 1–14, 57–70; offspring –, 509–522; parental –, 1169–1181; territory –, 902–914
Queen Maud Gulf Bird Sanctuary, 313–318
Quelea, 1271–1280
Quiscalus quiscula, 1130–1148

Rabosky, Daniel L., review by, 371–373
radiotelemetry, 36–49
Rail, 26–35; Black, 26–35; Clapper, 26–35; King, 26–35; Virginia, 26–35; Yellow, 26–35
rainfall, 571–592
rainforest, 108–120, 783–792, 1064–1069
Rallus elegans, 26–35
R. limicola, 26–35
R. l. levipes, 26–35
R. l. obsoletus, 26–35
R. l. yumanensis, 26–35
Ramphastidae, 1055–1063
Ramphastos, 1191–1209
R. ambiguus, 1055–1063, 1191–1209
R. brevis, 1191–1209
R. dicolorus, 1191–1209
R. sulfuratus, 1191–1209
R. s. brevicarinatus, 1191–1209
R. s. sulfuratus, 1191–1209
R. swainsonii, 1191–1209
R. toco, 1055–1063, 1191–1209
R. t. toco, 1191–1209
R. tucanus, 1191–1209
R. t. cuvieri, 1191–1209
R. t. tucanus, 1191–1209
R. vitellinus, 1055–1063, 1191–1209
R. v. ariel, 1191–1209
R. v. citreolaemus, 1191–1209
R. v. culminatus, 1191–1209
R. v. vitellinus, 1191–1209
Randia, 1271–1280
ranging behavior, 1249–1260
Raphus cucullatus, 1003–1004
Rappole, John H., et al., A new species of scimitar-babbler (Timaliidae: *Jabouilleia*) from the sub-Himalayan region of Myanmar, 1064–1069
raptor, 108–120
ratites, 1049–1054
Raven, Common, 36–49
Razorbill, 57–70, 773–782, 887–901
recognition, 530–543; enemy –, 530–543; species –, 689–700
Redhead, 1130–1148
Redstart, American, 71–81, 319–328, 733–746, 793–802, 1037–1048, 1070–1085, 1130–1148; Black, 71–81
Redwing, 733–746
Reed, Austin, see Menu, Stéphane, et al.
Rees, Eileen C., In Memoriam: Janet Kear, 1933–2004, 1293–1294
Regulus, 733–746
R. calendula, 71–81, 544–557, 689–700, 938–948, 1070–1085, 1130–1148
R. non cristatus, 706–707
R. satrapa, 1070–1085
regurgitations, 1182–1190
Remiz, 121–143
R. pendulinus, 121–143
Remsen, J. V., Jr., Pattern, process, and rigor meet classification, 403–413
renest, 306–312
renesting, 280–291, 902–914; – behavior, 1000–1001; – probability, 1001–1003
Renfrew, Rosalind B., et al., Edge avoidance by nesting grassland birds: A futile strategy in a fragmented landscape, 618–636
Renjifo, Luis Miguel, see Cuervo, Andrés M., et al.
Renner, Swen C., see Rappole, John H., et al.
repertoire, – size, 175–186; song –, 82–93, 175–186; duet –, 175–186; vocal –, 121–143
reproductive, – behavior, 347–348; – effort, 902–914; – investment, 509–522, 1169–1181; – isolation, 403–413; – output, 246–254, 306–312, 829–834; – physiology, 255–267; – success, 205–221, 571–592, 747–758, 759–772, 843–852, 1086–1096, 1169–1181, 1225–1238; – success, male, 558–565
reproductively isolated, 706–707; – demes, 803–818
resource, – availability, 544–557, 1281–1290; – defense, 1097–1111; – partitioning, 187–204; – exploitation, 689–700
retrospective research, 990–994
return rates, 329–337
Reviewers for *The Auk*, 2005, 1312–1313
Revillagigedo Archipelago, 50–56
Rhamphastidae, 1049–1054
Rheinwald, Goetz, see Stokke, Bård G., et al.
Rhinocrypta lanceolata, 445–463, 673–683
Rhinocryptidae, 445–463, 673–683
Rhopoterpe, 673–683
Rhymer, Judith M., et al., Phylogeography of the American Woodcock (*Scolopax minor*): Are management units based on band recovery data reflected in genetically based management units?, 1149–1160
Ribic, Christine A., see Renfrew, Rosalind B., et al.
Rice, Nathan H., Phylogenetic relationships of antpitta genera (Passeriformes: Formicariidae), 673–683
Rich, T. D., et al., Partners in Flight North American Landbird Conservation Plan (rev.), 373–375
ridges, 497–508
Rimator, 1064–1069
R. danjoui, 1064–1069
R. malacoptilus, 1064–1069
Rissa, 268–279
R. tridactyla, 268–279
Roadrunner, Greater, 523–529
Robbins, Chandler S., review by, 1305–1307
Robbins, Chandler S., review by, 375–377
Robbins, Chandler S., see Dunn, Erica H., et al.
Robin, American, 544–557, 1070–1085, 1130–1148; Rufous Bush, 1169–1181

Robinson, W. Douglas, et al., Are artificial bird nests effective surrogates for estimating predation on real bird nests? A test with tropical birds, 843–852
Rocque, Deborah A., and Winker, Kevin, Use of bird collections in contaminant and stable-isotope studies, 990–994
Rodríguez G., Malinalli, see Herrera M., L. Gerardo, et al.
Rohwer, Sievert, see Kulikova, Irina V., et al.
Roller, 1049–1054; Abyssinian, 1271–1280; Blue-throated, 1271–1280
Rosenberg, K. V., see Rich, T. D., et al.
Rosenberg, Kenneth V., see Dunn, Erica H., et al.
Rosivall, Balázs, et al., Food allocation in Collared Flycatcher (*Ficedula albicollis*) broods: Do rules change with the age of nestlings?, 1112–1122
Rosivall, Balázs, see Hargitai, Rita, et al.
Rothschild, Miriam, see van Emden, Helmet F., and –
Rothstein, Stephen I., see Anderson, Kurt E., et al.
Rothstein, Stephen I., see Hosoi, S. Aki, et al.
Rupelramphastoides, 1055–1063
R. knopfi, 1055–1063
Rupicola rupicola, 793–802
Rustay, C. M., see Rich, T. D., et al.

Sabrewings, Long-tailed, 593–607; Wedge-tailed, 593–607
Sæther, Bernt-Erik, see Stokke, Bård G., et al.
Saito, Daichi S., see Hamao, Shoji, and –
salinity, 523–529
salt, – glands, 523–529; – marsh, 94–107; – stress, 523–529
Salt, George William, 1919–1999, In Memoriam, 354
saltcedar, 1261–1270
Sampieri, Sara J., and Clotfelter, Ethan D., Temporal and geographic patterns of age-specific gull plumage in relation to potential exposure to polychlorinated biphenyls, 915–921
Sandpiper, 1291–1292; Green, 1291–1292; Spotted, 1130–1148; Wood, 1291–1292
Sapphire, Blue-headed, 1026–1031; Humboldt's, 1026–1031
Sapsucker, Red-naped, 1070–1085; Yellow-bellied, 938–948, 938–948
Saroglossa, 1271–1280
Sasia, 1055–1063
S. africana, 1055–1063
Sattler, Gene D., see Bronson, C. L., et al.
Sauer, John R., see Dunn, Erica H., et al.
Sayornis phoebe, 1130–1148
Scaup, Lesser, 1130–1148
Schwagmeyer, P. L., see Bartlett, Terri L., et al.
Schwalbenkatastrophe, 637–647
Scimitar-Babbler, 1064–1069; **Naung Mung**, 1064–1069; Short-tailed, 1064–1069
Scolopax, 1249–1260
S. minor, 15–25, 1149–1160
S. rusticola, 1149–1160
Scrub-Jay, Florida, 464–478, 571–592; Western, 464–478, 571–592
scutellation, 673–683
Scytalopus, 445–463
S. aff. *bolivianus*, 445–463
S. affinis, 445–463
S. argentifrons, 445–463
S. atratus, 445–463
S. a. confusus, 445–463
S. canus opacus, 445–463
S. chocoensis, 445–463
S. femoralis, 445–463
S. fuscus, 445–463
S. griseicollis, 445–463
S. latebricola, 445–463
S. latrans, 445–463
S. l. latrans, 445–463
S. magellanicus, 445–463
S. micropterus, 445–463
S. parkeri, 445–463
S. parvirostris, 445–463
S. robbinsi, 445–463
S. sanctaemartae, 445–463
S. schulenbergi, 445–463
S. simonsi, 445–463
S. spillmanni, 445–463
S. stilesi, 445–463
S. subcinereus, 445–463
S. superciliaris, 445–463
S. vicinior, 445–463
S. zimmeri, 445–463
sea duck, 313–318
seabirds, 15–25, 773–782, 887–901, 994–999, 1182–1190
Sealy, S. G., Letters to *The Auk*, 347
Sealy, Spencer G., et al., In Memoriam: Roger Malcolm Evans, 1935–1998, 351–353
Sealy, Spencer G., see Murphy-Klassen, Heather M., et al.
search image, 36–49
Searcy, William A., see Nowicki, Stephen, and –
seasonal interactions, 1037–1048
Seed-cracker, Black-bellied, 1123–1129
Seedeater, Tumaco, 389–402
Seiurus aurocapilla, 280–291, 793–802, 938–948
S. motacilla, 938–948
S. noveboracensis, 319–328, 938–948
Selasphorus rufus, 187–204, 872–886, 938–948, 1070–1085
S. sasin, 872–886
selection, ecological –, 689–700; habitat –, 1097–1111, 1249–1260; sexual –, 57–70, 82–93, 689–700, 872–886, 1161–1168; site –, 1249–1260
Selenidera culik, 1055–1063
S. maculirostris, 1055–1063
S. reinwardtii, 1191–1209
sensory, – bias, 1–14; – drive, 689–700
Sephanoides fernandensis, 593–607

Serinus canaria, 1–14
Setophaga ruticilla, 71–81, 319–328, 733–746, 793–802, 1037–1048, 1070–1085, 1130–1148
sex, – differences, 175–186, 1239–1248; – ratio, 1161–1168
sexing, – birds, 347–348; – criteria, 347–348
sex-limited inheritance, 1123–1129
sex-specific, – behavior, 347–348; – distributions, 938–948
sexual, – dimorphism, 187–204, 347–348; – segregation, 938–948; – selection, 57–70, 82–93, 689–700, 872–886, 1161–1168; – size dimorphism, 922–937
sexually dimorphic plumage, 835–842
Shaffer, Terry L., see Grant, Todd A., et al.
Shag, European, 773–782
Shearwater, 608–617; Cory's, 255–267
Sheets, H. David, see Morris, Sara R., et al.
Sheldon, Frederick H., see Gill, Frank B., et al.
Shikra, 108–120
shorebirds, 15–25, 479–496, 922–937, 1149–1160, 1249–1260
Shoveler, Northern, 1130–1148
Shrike, Loggerhead, 1037–1048
Shriver, W. Gregory, et al., Concordance between morphological and molecular markers in assessing hybridization between Sharp-tailed Sparrows in New England, 94–107
Shugar, Martin A., see Currie, Phillip J., et al.
Shwe, Nay Myo, see Rappole, John H., et al.
Sialia currucoides, 1130–1148
S. sialis, 1086–1096
sibling species, 706–707, 1291–1292
Silveira, Luís Fábio, et al., A new species of *Aratinga* parakeet (Psittaciformes: Psittacidae) from Brazil, with taxonomic remarks on the *Aratinga solstitialis* complex, 292–305
Simons, Theodore R., see Farnsworth, George L., and –
Siskin, Pine, 1070–1085
site, – fidelity, 329–337; – selection, 1249–1260; stopover –, 319–328, 1281–1290; subarctic stopover –, 544–557
Sitta canadensis, 1070–1085
S. pusilla, 1026–1031
Sittiparus, 121–143
S. varius, 747–758
skeletal characters, 673–683
skeletons, 971–979, 1055–1063
skewed sex ratios, 938–948
Skua, 1182–1190, Great, 1037–1048
Skutch, Alexander F., 1904–2004, In Memoriam, 708–710
Slagsvold, Tore, In Memoriam: Svein Haftorn, 1925–2003, 358
Slikas, Beth, see Gill, Frank B., et al.
Smith, Kimberly G., 100 Years Ago in the American Ornithologists' Union, 1310–1311
Smith, Kimberly G., see Dunn, Erica H., et al.
Snipe, Common, 1149–1160; Wilson's, 1130–1148
social, – adaptation, 803–818; – attraction, 689–700; – dominance, 187–204
Socorro Island, 50–56
soft-part colors, 673–683
Sogge, Mark K., see Owen, Jennifer C., et al.
Solitaire, 1003–1004; Townsend's, 1070–1085
Somateria mollissima, 523–529
S. spectabilis, 313–318
song, 1–14, 82–93, 175–186, 231–245; adult –, 853–871; – dialects, 231–245, 593–607, 803–818; juvenile –, 853–871; – learning, 1–14, 231–245, 853–871; – repertoire, 82–93, 175–186; solo –, 175–186; – structure, 593–607
songbirds, 36–49, 82–93, 121–143, 280–291, 280–291, 319–328, 803–818, 843–852, 853–871, 902–914, 1055–1063
Sora, 26–35, 1130–1148
Sorenson, Michael D., see Kulikova, Irina V., et al.
sound, 984–987; – transmission, 231–245
Sparrow, 94–107, 389–402, 403–413; American Tree, 544–557, 1130–1148; Baird's, 349–350; Brewer's, 571–592, 1070–1085; Chipping, 1070–1085; Clay-colored, 661–672, 701–705, 938–948, 1130–1148; Field, 530–543; Fox, 530–543, 544–557, 1070–1085, 1130–1148; Gambel's White-crowned, 1070–1085; Golden-crowned, 1070–1085; Grasshopper, 94–107, 618–636; Harris's, 1130–1148; Henslow's, 618–636, 661–672; House, 57–70, 835–842, 1086–1096; Ipswich, 403–413; Lark, 1070–1085; Lincoln's, 544–557, 938–948, 1070–1085; Nelson's Sharp-tailed, 94–107; Nuttall's White-crowned, 853–871; Rock, 1086–1096; Rufous-collared, 231–245; Rufous-crowned, 571–592; Sage, 571–592; Saltmarsh Sharp-tailed, 94–107; Savannah, 403–413, 544–557, 618–636, 701–705; Song, 1–14, 82–93, 280–291, 403–413, 571–592, 648–660, 853–871, 1130–1148; Swamp, 1–14; Vesper, 661–672, 701–705, 1070–1085, 1130–1148; White-crowned, 82–93, 544–557, 803–818, 938–948, 1070–1085, 1130–1148; White-throated, 71–81, 1130–1148
Sparrowhawk, Besra, 108–120
spatial, – memory, 608–617, 819–828; – patterns, 338–346
speciation, 268–279, 414–432, 433–444, 706–707, 759–772; recent –, 684–688
species, – composition, 1097–1111; biological – concept, 50–56, 403–413, 433–444; – concepts, 94–107, 292–305; – limits, 433–444, 445–463, 1191–1209; – recognition, 689–700; – richness, 338–346
specimen salvage, 966–971
spectophotometry, reflectance –, 793–802
spectrograms, 733–746, 853–871, 984–987
speculation, 347
Sphenisciformes, 971–979
Sphyrapicus nuchalis, 1070–1085

S. varius, 938–948, 1055–1063
Spilornis cheela, 108–120
Spinebill, Eastern, 1097–1111
Spiza americana, 82–93, 648–660, 661–672
Spizella arborea, 544–557, 1130–1148
S. breweri, 571–592, 1070–1085
S. pallida, 661–672, 701–705, 938–948, 1130–1148
S. passerina, 1070–1085
S. pusilla, 530–543
Sporophila insulata, 389–402
spring, – arrivals, 1130–1148; – fatter hypothesis, 544–557
squirrels, 108–120
Sri Lanka, 108–120
stable isotopes, 990–994, 1037–1048
stable-isotope analysis, 783–792
Stacey, Peter B., see Arsenault, David P., et al.
Starling, 608–617, 1271–1280; European, 1–14, 222–230, 479–496, 835–842; Rose-coloured, 1037–1048; Wattled, 1271–1280
Stellula calliope, 1070–1085
Stepney, Philip H. R., 1947–2004, In Memoriam, 711
Sterna, 268–279
S. eurygnatha, 268–279
S. forsteri, 1130–1148
Sternidae, 268–279
Sterocorarius skua, 1037–1048
Stiles, F. Gary, et al., Wing morphology and flight behavior of some North American hummingbird species, 872–886
Stiles, F. Gary, In Memoriam: Alexander F. Skutch, 1904–2004, 708–710
Stokke, Bård G., et al., Weather in the breeding area and during migration affects the demography of a small long-distance passerine migrant, 637–647
stomach samples, 1182–1190
stopover, – duration, 319–328; – ecology, 544–557; – sites, 319–328, 1281–1290
Storer, Robert W., A possible connection between crop milk and the maximum size attainable by flightless pigeons, 1003–1004
storks, 608–617
Storm-Petrel, Swinhoe's, 733–746
Streptopelia, 1271–1280
S. decaocto, 530–543
Strigiformes, 971–979, 1130–1148
Strix occidentalis caurina, 329–337
Sturnella, 618–636
S. magna, 618–636, 648–660, 661–672
S. neglecta, 82–93, 661–672, 1130–1148
Sturnus roseus, 1037–1048
S. vulgaris, 1–14, 222–230, 479–496, 835–842
Stutchbury, Bridget J. M., see Gill, Sharon A., and –
Styrsky, Jennifer Nesbitt, see Robinson, W. Douglas, et al.
Sub-Himalayan region, 1064–1069
subird-asities, 1271–1280
suboscine, 403–413, 433–444, 445–463, 673–683
subpopulations, 329–337
subspecies, 433–444
Sunbird, 187–204; Orange-tufted, 231–245
survey, 15–25; – methods, 26–35
survival, 479–496, 523–529, 637–647, 773–782, 829–834; juvenile –, 205–221; nest –, 618–636, 661–672; postfledging –, 479–496; time-specific –, 661–672
survivorship, 1037–1048
Sutherland, William J., see Gillings, Simon, et al.
Sutter, Robert D., see Mehlman, David W., et al.
Svärdson, Gunnar, 1914–2004, In Memoriam, 1011–1012
Swallow, 608–617, 733–746; Barn, 509–522, 637–647, 1130–1148; Cliff, 121–143, 637–647, 1225–1238; Mangrove, 1026–1031; Tree, 57–70, 246–254, 915–921, 1130–1148, 1225–1238
Swan, Tundra, 1130–1148
Swanson, David L., see Carlisle, Jay D., et al.
Sweet, Paul R., see Rappole, John H., et al.
Swift, Chapman's, 389–402
Sylvia atricapilla, 121–143, 530–543, 733–746
S. borin, 608–617, 1037–1048
S. melanocephala, 608–617
Sylviidae, 733–746
Sylviparus, 121–143
S. modestus, 121–143
sympatric distributions, 158–174
sympatry, 158–174
systematic paleontology, 1055–1063
systematics, 268–279, 292–305, 389–402, 414–432, 433–444, 445–463, 987–990, 1037–1048
Szigeti, Beáta, see Hargitai, Rita, et al.
Szöllősi, Eszter, see Rosivall, Balázs, et al.
Szölőlsi, Eszter, see Hargitai, Rita, et al.

Tachycineta albilinea, 1026–1031
T. bicolor, 57–70, 246–254, 915–921, 1130–1148, 1225–1238
Tadorna tadornoides, 523–529
Taenopygia guttata, 1–14, 255–267, 509–522, 1112–1122, 1239–1248
tallgrass prairie, 648–660
tamarisk, 1261–1270
Tanager, 733–746, 938–948; Summer, 938–948; Western, 938–948, 1070–1085
Tapaculo, 673–683; **Stiles's**, 445–463
taxonomic status, 292–305, 433–444, 684–688
taxonomy, 121–143, 158–174, 445–463, 673–683, 966–971
Teal, Blue-winged, 1130–1148; Green-winged, 1130–1148
Temeles, Ethan J., et al., Foraging and territory economics of sexually dimorphic Purple-throated Caribs (*Eulampis jugularis*) on three *Heliconia* morphs, 187–204
temperate migrants, 1070–1085

temperature, 571–592, 1130–1148
terminal investment hypothesis, 509–522
Tern, 15–25; Black, 1130–1148; Cayenne, 268–279; Forster's, 1130–1148; White, 1026–1031
Terpsiphone, 1271–1280
territoriality, 71–81, 187–204
territories, 497–508, 593–607
territory, – density, 497–508; – quality, 902–914
Thalassarche chrysostoma, 1182–1190
T. melanophris, 1182–1190
Thamnocharis, 673–683
Thamnophilidae, 403–413, 414–432, 433–444, 673–683
Thamnophilus aethiops, 414–432
T. atrinucha, 433–444
T. caerulescens, 403–413, 414–432, 433–444
T. c. aspersiventer, 414–432, 433–444
T. c. connectens, 403–413, 414–432, 433–444
T. c. dinellii, 414–432, 433–444
T. c. paraguayensis, 414–432, 433–444
T. punctatus, 292–305
T. unicolor, 414–432
Thectocercus, 292–305
Thévenot, Michel, et al., The Birds of Morocco (rev.), 1303–1305
Thompson, Christopher W., see Lance, Monique M., and –
Thornbill, Buff-rumped, 1097–1111
Thrasher, Brown, 1130–1148
Thrush, 733–746, 938–948, 1271–1280; Gray-cheeked, 544–557, 733–746; Hermit, 689–700, 938–948, 1070–1085, 1130–1148; Song, 530–543; Swainson's, 94–107, 544–557, 733–746, 938–948, 938–948, 1037–1048, 1070–1085; Wood, 280–291, 306–312
Thryomanes, 50–56
T. bewickii, 50–56
T. insularis, 50–56
T. sissonii, 50–56
T. (*Troglodytes*) *sissonii*, 50–56
Thryorchilus browni, 50–56
Thryothorus, 175–186
T. (*ludovicianus*) *albinucha*, 50–56
T. fasciatoventris, 175–186
T. f. albigularis, 389–402
T. felix, 175–186
T. leucotis, 175–186, 1169–1181
T. ludovicianus, 50–56, 175–186
T. modestus, 175–186
T. nicefori, 175–186
T. nigricapillus, 175–186
T. pleurostictus, 175–186
T. rufalbus, 175–186
T. rutilis, 175–186
T. sinaloa, 175–186
T. sissonii, 50–56
Timaliidae, 108–120
Timewell, Christopher A. R., see Mac Nally, Ralph, and –
timing of breeding, 829–834, 902–914
Tinamiformes, 971–979
tinamou, 971–979
Tipton, John R., see Gammon, David E., et al.
Tit, 121–143, 1225–1238; Azure, 747–758; Black-crested, 747–758; Bearded, 689–700; Blue, 57–70, 509–522, 558–565, 747–758, 793–802, 1086–1096; Coal, 747–758; Great, 57–70, 71–81, 231–245, 246–254, 558–565, 747–758, 793–802, 829–834, 902–914, 1130–1148; Siberian, 819–828; Turkestan, 747–758; Varied, 747–758; Willow, 108–120, 747–758, 819–828; Yellow-breasted, 747–758
Titmouse, 121–143, 747–758, 1023; Black-crested, 121–143, 747–758; Bridled, 121–143; Juniper, 121–143, 747–758; Oak, 121–143, 747–758; Tufted, 108–120, 121–143, 747–758
Tockus nasutus, 1271–1280
Todiramphus chloris, 1271–1280
Török, János, see Hargitai, Rita, et al.
Török, János, see Rosivall, Balázs, et al.
Tóth, László, see Hargitai, Rita, et al.
Toucan, 1049–1054, 1055–1063, 1191–1209; Ariel, 1191–1209; Black-mandibled, 1191–1209; Channel-billed, 1191–1209; Chestnut-mandibled, 1191–1209; Choco, 1191–1209; Cuvier's, 1191–1209; Hooded Mountain, 1191–1209; Keel-billed, 1191–1209; Red-billed, 1191–1209; Red-breasted, 1191–1209; Toco, 1191–1209; Yellow-ridged, 1191–1209
Toucanet, Emerald, 1191–1209; Golden-collared, 1191–1209; Saffron, 1191–1209
Towhee, 403–413; Spotted, 1070–1085
toxicology, 255–267
Toxostoma rufum, 1130–1148
tracheophone, 673–683
Trachyphonus, 1055–1063
T. margaritatus, 1055–1063
trapline-forage, 187–204
Traylor, Joshua J., see Kellett, Dana K., et al.
Trimble, Jeremiah, see Causey, Douglas, and –
Tringa flavipes, 1130–1148
T. glareola, 1291–1292
T. melanoleuca, 1130–1148
T. ochropus, 1291–1292
Trochilidae, 872–886, 1055–1063
Trochilus humboldtii, 1026–1031
Troglodytes, 50–56
T. aedon, 50–56, 689–700, 1070–1085, 1130–1148
T. beani, 50–56
T. brunneicollis, 50–56
T. [= *musculus*] *inquietus*, 50–56
T. insularis, 50–56
T. musculus, 50–56
T. ochraceus, 50–56
T. rufociliatus, 50–56
T. rufulus, 50–56
T. sissonii, 50–56

T. solstitialis, 50–56
T. tanneri, 50–56
T. troglodytes, 50–56
T. (*Nannus*) *troglodytes*, 50–56
Troglodytidae, 50–56, 1026–1031
Trogon, Malabar, 108–120
tropical, – birds, 783–792; – forest, 843–852
trypanosomes, 558–565
tubenoses, 1055–1063
tundra, 144–157
Turdoides rufescens, 108–120
Turdus, 733–746
T. iliacus, 733–746
T. merula, 530–543, 558–565, 733–746
T. migratorius, 544–557, 1070–1085, 1130–1148
T. philomelos, 530–543
T. pilaris, 733–746
Turkey, Wild, 618–636
Turnstone, Ruddy, 1149–1160
type locality, 347–348
Tyrannidae, 733–746, 938–948, 1055–1063
tyrannoids, 673–683
Tyrannus tyrannus, 1130–1148
T. verticalis, 1130–1148
Tyto alba, 733–746

Ulfstrand, Staffan, In Memoriam: Gunnar Svärdson, 1914–2004, 1011–1012
ultraviolet, 793–802; – degradation, 793–802
understory, 1064–1069
Underwood, Todd J., see Murphy-Klassen, Heather M., et al.
Upucerthia, 158–174
U. ruficauda, 158–174
urban, – acoustics, 231–245; – ecology, 205–221; – sprawl, 205–221
Urban, Emil K., review by, 1303–1305
Uria aalge, 773–782, 887–901
U. lomvia, 887–901

van Emden, Helmet F., and Rothschild, Miriam, Insect and Bird Interactions (rev.), 1019–1020
Vanellus vanellus, 1249–1260
Vangidae, 1271–1280
variation, clinal –, 414–432, 433–444; geographic –, 403–413, 648–660, 803–818, 853–871; interclutch –, 509–522; intraclutch –, 509–522; microgeographic song –, 593–607; mitochondrial –, 414–432
Veery, 733–746
vegetative concealment, 648–660
Vehrencamp, Sandra L., see Mennill, Daniel J., and –
Veit, Richard R., review by, 1307–1308
Vermivora celata, 544–557, 938–948, 1070–1085
V. peregrina, 1037–1048
V. ruficapilla, 938–948, 1070–1085
Verner, Jared, 361–362
Vernon, Rae, see Thévenot, Michel, et al.
vicariant events, 158–174
Vickers-Rich, Patricia, see Murray, Peter F., and –
Vickery, Peter D., see Shriver, W. Gregory, et al.
video-monitored nests, 701–705
viduine, 403–413
Vigallon, Stacey M., and Marzluff, John M., Is nest predation by Steller's Jays (*Cyanocitta stelleri*) incidental or the result of a specialized search strategy?, 36–49
Vireo, 733–746, 938–948; Bell's, 306–312, 938–948; Black-capped, 280–291; Blue-headed, 280–291, 938–948; Cassin's, 1070–1085; Puerto Rican, 280–291; Red-eyed, 280–291, 319–328, 1070–1085; Warbling, 280–291, 1070–1085
Vireo atricapilla, 280–291
V. bellii, 306–312, 938–948
V. cassinii, 1070–1085
V. gilvus swainsonii, 1070–1085
V. gilvus, 280–291, 1070–1085
V. olivaceus, 280–291, 319–328, 1070–1085
V. solitarius, 280–291, 938–948
Vireonidae, 733–746
vocal, – behavior, 175–186; – dialects, 231–245; – performance, 1–14
vocalization, 403–413, 414–432, 433–444, 445–463, 593–607, 673–683, 733–746, 853–871, 1191–1209; – probability, 26–35
Voous, Karel Hendrik, 1920–2002, In Memoriam, 355–356
vouchers, digital –, 979–984
Vuilleumier, François, Dean of American ornithologists: The multiple legacies of Frank M. Chapman of the American Museum of Natural History, 389–402
Vuilleumier, François, In Memoriam: Karel Hendrik Voous, 1920–2002, 355–356
Vuilleumier, François, see Cheviron, Z. A., et al.
Vulture, Turkey, 389–402

wader, 15–25
wading birds, 255–267
Wagtail, 733–746; Black-backed, 1026–1031; Madagascar, 1271–1280
Waldronia milleri, 389–402
Walters, Michael, A Concise History of Ornithology (rev.), 716–718
Warbler, 608–617, 733–746, 938–948, 1024–1025, 1271–1280; Aquatic, 558–565; Black-and-white, 793–802, 938–948; Black-browed Reed, 1086–1096; Black-throated Blue, 71–81, 306–312, 571–592, 793–802, 902–914, 1000–1001, 1001–1003, 1037–1048, 1086–1096; Black-throated Gray, 938–948; Black-throated Green, 793–802, 938–948; Blackpoll, 544–557, 689–700, 733–746; Canada, 793–802; Cerulean, 497–508; Chestnut-sided, 793–802; Clamorous Reed, 558–565; Dusky, 71–81; Garden, 608–617, 1037–1048; Grasshopper, 1291–1292; Great Reed, 1–14,

530–543, 558–565, 1086–1096; Hermit, 938–948, 1070–1085; Hooded, 902–914, 938–948, 1086–1096; Kentucky, 938–948, 1037–1048; Kirtland's, 689–700; MacGillivray's, 938–948, 1070–1085; Madagascar Swamp, 1271–1280; Magnolia, 71–81, 793–802, 938–948, 1070–1085; Moustached, 835–842; Nashville, 938–948, 1070–1085; Orange-crowned, 544–557, 938–948, 1070–1085; Palm, 1130–1148; Prairie, 280–291, 689–700; Reed, 530–543, 558–565, 835–842, 1291–1292; River, 733–746; Sardinian, 608–617; Sedge, 558–565; Sevi's, 1291–1292; Swainson's, 689–700; Tennessee, 1037–1048; Townsend's, 938–948, 1070–1085; Willow, 530–543, 1291–1292; Wilson's, 938–948, 1070–1085, 1130–1148; Wood, 1291–1292; Worm-eating, 280–291; Yellow, 94–107, 530–543, 544–557, 938–948, 1070–1085, 1130–1148; Yellow-rumped, 71–81, 544–557, 733–746, 793–802, 938–948, 1037–1048, 1070–1085, 1130–1148
Warnell, Daniel B., see Cooper, Robert J., and –
Washington, 36–49, 887–901
waterbird, 15–25, 1049–1054
waterfowl, 144–157, 313–318, 523–529, 618–636, 949–965, 1049–1054, 1130–1148
Waterthrush, Louisiana, 938–948; Northern, 319–328, 938–948
Wattlebird, Red, 1097–1111
Waxwing, Cedar, 1070–1085
Weakland, Cathy A., and Wood, Petra Bohall, Cerulean Warbler (*Dendroica cerulea*) microhabitat and landscape-level habitat characteristics in southern West Virginia, 497–508
weather, 571–592, 637–647, 1130–1148, 1225–1238
weaverbird, 57–70
weavers, 57–70
Weckstein, Jason D., Molecular phylogenetics of the *Ramphastos* toucans: Implications for the evolution of morphology, vocalizations, and coloration, 1191–1209
Wendt, J. S., see Rich, T. D., et al.
wetlands, 26–35, 255–267
Whitaker, Justine M., see Cristol, Daniel A., et al.
Whiteman, Noah K., see Bollmer, Jennifer L., et al.
Whittingham, Linda A., see Nooker, Jacqueline K., et al.
widows, 57–70
Wieczorek, John, see Peterson, A. Townsend, et al.
Wigeon, American, 1130–1148
Wiley, James W., see Keith, Allan R., et al.
Will, T. C., see Rich, T. D., et al.
Willet, 1130–1148
willow-wrens, 706–707, 1291–1292
Wilsonia canadensis, 793–802
W. citrina, 902–914, 938–948, 1086–1096
wing disc loading, 872–886; – loading, 872–886
Wink, Michael, see Dyrcz, Andrzej, et al.
Winker, Kevin, Bird collections: Development and use of a scientific resource, 966–971
Winker, Kevin, see Benson, Anna-Marie, and –
Winker, Kevin, see Kulikova, Irina V., et al.
Winker, Kevin, see Rocque, Deborah A., and –
Winker, Kevin, Sibling species were first recognized by William Derham (1718), 706–707
winter ecology, 71–81; – site-fidelity, 608–617
Wisonia pusilla, 544–557, 938–948, 1070–1085, 1130–1148
Witek, Magdalena, see Dubiec, Anna, et al.
Withey, John C., and Marzluff, John M., Dispersal by juvenile American Crows (*Corvus brachyrhynchos*) influences population dynamics across a gradient of urbanization, 205–221
Wood, Petra Bohall, see Weakland, Cathy A., and –
Woodcock, 1249–1260; American, 15–25, 1149–1160; Eurasian, 1149–1160
Woodpecker, 733–746, 938–948, 1049–1054, 1055–1063, 1191–1209; Beautiful, 1026–1031; Downy, 108–120; Ivory-billed, 389–402, 984–987; Red-bellied, 558–565
Wood-Pewee, Eastern, 1070–1085; Western, 82–93, 1070–1085
Woodrey, Mark, see Mehlman, David W., et al.
wood-warblers, 71–81, 733–746, 747–758, 793–802, 1037–1048
Wood-Wren, White-breasted, 50–56
Wrege, Peter H., and Emlen, Stephen T., Sexing criteria, accuracy, and statistical inference — A reply, 348–349.
Wren, Banded, 175–186; Bay, 175–186; Bewick's, 50–56; Black-bellied, 175–186, 389–402; Brown-throated, 50–56; Buff-breasted, 175–186, 1169–1181; Carolina, 50–56, 175–186; Clarion, 50–56; Cozumel, 50–56; Happy, 175–186; House, 50–56, 689–700, 1070–1085, 1130–1148; Long-billed Marsh, 389–402; Marsh, 26–35, 1130–1148; Mountain, 50–56; Niceforo's, 175–186; Northern House, 50–56; Ochraceous, 50–56; Plain, 175–186; Rufous-and-white, 175–186; Rufous-breasted, 175–186; Rufous-browed, 50–56; Sedge, 50–56; Sinaloa, 175–186; Socorro, 50–56; Southern House, 50–56; Tepui, 50–56; Timberline, 50–56; White-browed, 50–56; Winter, 50–56
Wright, Joanna L., see Currie, Phillip J., et al.
Wright, Jonathan, see Leader, Noam, et al.
wrynecks, 1055–1063

Xanthocephalus xanthocephalus, 1130–1148
Xavier, José C., et al., Boluses: An effective method for assessing the proportions of cephalopods in the diet of albatrosses, 1182–1190

Yellowlegs, Greater, 1130–1148; Lesser, 1130–1148
Yellowthroat, Common, 793–802, 1070–1085, 1130–1148

yelpers, smooth-billed, 1191–1209
Yom-Tov, Yoram, see Leader, Noam, et al.
young, 479–496

Zenaida macroura, 1130–1148
Zhuravlev, Yuri N., see Kulikova, Irina V., et al.
Ziegenfus, Charles W. S., see Keiser, Jeffrey T., et al.
Ziel, Heather L., see Rhymer, Judith M., et al.
Zonotrichia albicollis, 71–81, 1130–1148
Z. atricapilla, 1070–1085
Z. capensis, 231–245
Z. leucophrys, 82–93, 544–557, 803–818, 938–948, 1070–1085, 1130–1148
Z. l. gambeli, 1070–1085
Z. l. nuttalli, 853–871
Z. querula, 1130–1148
Zosterops, 1271–1280
zugunruhe, 733–746